Faculty of Electrical Engineering Czech Technical University in Prague

## Projections onto spaces of polynomials

Tommaso Russo (Joint work in progress with P. Hájek)

Workshop on Banach spaces and Banach lattices Madrid, Spain September 9–13, 2019

#### International Mobility of Researchers in CTU Project number: CZ.02.2.69/0.0/0.0/16\_027/0008465



EVROPSKÁ UNIE Evropské strukturální a investiční fondy Operační program Výzkum, vývoj a vzdělávání



- ▶ A Banach space X has the AP if for every compact set  $K \subseteq X$  and  $\varepsilon > 0$  there exists a finite-rank, bounded linear operator  $T: X \to X$  such that  $||Tx x|| < \varepsilon$  ( $x \in K$ );
- X has  $\lambda$ -BAP if additionally  $||T|| \leq \lambda$ .

#### Theorem (Godefroy and Kalton, 2003

A Banach space X has the  $\lambda$ -BAP if and only if  $\mathcal{F}(X)$  has the  $\lambda$ -BAP.

In particular,  $\mathcal{F}(\ell_2)$  has the MAP ( $\equiv$  1-BAP).

#### Problem (Godefroy)

Does  $Lip_0(\ell_2)$  have the AP?

- **Grothendieck (1955).** The AP passes from  $X^*$  to X
- Hence, this would be a stronger result.

- ▶ A Banach space X has the AP if for every compact set  $K \subseteq X$  and  $\varepsilon > 0$  there exists a finite-rank, bounded linear operator  $T: X \to X$  such that  $||Tx x|| < \varepsilon$  ( $x \in K$ );
- X has  $\lambda$ -BAP if additionally  $||T|| \leq \lambda$ .

## Theorem (Godefroy and Kalton, 2003)

A Banach space X has the  $\lambda$ -BAP if and only if  $\mathcal{F}(X)$  has the  $\lambda$ -BAP.

In particular,  $\mathcal{F}(\ell_2)$  has the MAP ( $\equiv$  1-BAP)

#### Problem (Godefroy)

Does  $Lip_0(\ell_2)$  have the AP?

- **Grothendieck (1955).** The AP passes from X\* to X
- Hence, this would be a stronger result.

- ▶ A Banach space X has the AP if for every compact set  $K \subseteq X$  and  $\varepsilon > 0$  there exists a finite-rank, bounded linear operator  $T: X \to X$  such that  $||Tx x|| < \varepsilon$  ( $x \in K$ );
- X has  $\lambda$ -BAP if additionally  $||T|| \leq \lambda$ .

## Theorem (Godefroy and Kalton, 2003)

A Banach space X has the  $\lambda$ -BAP if and only if  $\mathcal{F}(X)$  has the  $\lambda$ -BAP.

• In particular,  $\mathcal{F}(\ell_2)$  has the MAP ( $\equiv$  1-BAP).

#### Problem (Godefroy)

Does  $Lip_0(\ell_2)$  have the AP?

- Grothendieck (1955). The AP passes from X\* to X;
- Hence, this would be a stronger result.

- ▶ A Banach space X has the AP if for every compact set  $K \subseteq X$  and  $\varepsilon > 0$  there exists a finite-rank, bounded linear operator  $T: X \to X$  such that  $||Tx x|| < \varepsilon$  ( $x \in K$ );
- X has  $\lambda$ -BAP if additionally  $||T|| \leq \lambda$ .

## Theorem (Godefroy and Kalton, 2003)

A Banach space X has the  $\lambda$ -BAP if and only if  $\mathcal{F}(X)$  has the  $\lambda$ -BAP.

• In particular,  $\mathcal{F}(\ell_2)$  has the MAP ( $\equiv$  1-BAP).

#### Problem (Godefroy)

Does  $Lip_0(\ell_2)$  have the AP?

- **Grothendieck (1955).** The AP passes from X\* to X;
- Hence, this would be a stronger result.

- ▶ A Banach space X has the AP if for every compact set  $K \subseteq X$  and  $\varepsilon > 0$  there exists a finite-rank, bounded linear operator  $T: X \to X$  such that  $||Tx x|| < \varepsilon$  ( $x \in K$ );
- X has  $\lambda$ -BAP if additionally  $||T|| \leq \lambda$ .

#### Theorem (Godefroy and Kalton, 2003)

A Banach space X has the  $\lambda$ -BAP if and only if  $\mathcal{F}(X)$  has the  $\lambda$ -BAP.

• In particular,  $\mathcal{F}(\ell_2)$  has the MAP ( $\equiv$  1-BAP).

#### Problem (Godefroy)

Does  $Lip_0(\ell_2)$  have the AP?

Grothendieck (1955). The AP passes from X<sup>\*</sup> to X;

Hence, this would be a stronger result.

- ▶ A Banach space X has the AP if for every compact set  $K \subseteq X$  and  $\varepsilon > 0$  there exists a finite-rank, bounded linear operator  $T: X \to X$  such that  $||Tx x|| < \varepsilon$  ( $x \in K$ );
- X has  $\lambda$ -BAP if additionally  $||T|| \leq \lambda$ .

#### Theorem (Godefroy and Kalton, 2003)

A Banach space X has the  $\lambda$ -BAP if and only if  $\mathcal{F}(X)$  has the  $\lambda$ -BAP.

• In particular,  $\mathcal{F}(\ell_2)$  has the MAP ( $\equiv$  1-BAP).

#### Problem (Godefroy)

Does  $Lip_0(\ell_2)$  have the AP?

- Grothendieck (1955). The AP passes from X<sup>\*</sup> to X;
- Hence, this would be a stronger result.

# 2

## For a Banach space X, $\mathcal{P}(^2X)$ is a subspace of $Lip_0(X)$ .

- ▶  $\mathcal{P}(^2X)$  is the Banach space of bounded 2-, on ogneous polynomials on X.
  - ▶  $P \in \mathcal{P}(^2X)$  if there is a bounded bilinear matrix  $X \to \mathbb{R}$  su that P(x) = M(x, x);
  - $||P||_{\mathcal{P}} = \sup_{x \in B_X} |P(x)|.$
- But polynomials are **not** Lipschitz functions!
- However, they are Lipschitz on the unit ball.
  - Therefore,  $\mathcal{P}(^2X)$  is a natural subspace of *Li*
  - Moreover,  $\|\cdot\|_{\mathcal{P}}$  is equivalent to  $\|\cdot\|_{Lip}$ .
- Consequently,  $\mathcal{P}(^2X)$  is naturally isomorphic t Lip<sub>0</sub>(B<sub>X</sub>), via the restriction map



For a Banach space X,  $\mathcal{P}(^2X)$  is a subspace of  $Lip_0(X)$ .

- $\mathcal{P}(^2X)$  is the Banach space of bounded 2-homogeneous polynomials on X.
  - ▶  $P \in \mathcal{P}(^2X)$  if there is a bounded bilinear mathematical that P(x) = M(x, x);
  - $||P||_{\mathcal{P}} = \sup_{x \in B_X} |P(x)|.$

But polynomials are **not** Lipschitz functions!

- However, they are Lipschitz on the unit ball.
  - Therefore,  $\mathcal{P}(^2X)$  is a natural subspace of *Li*
  - Moreover,  $\|\cdot\|_{\mathcal{P}}$  is equivalent to  $\|\cdot\|_{Lip}$ .

Consequently,  $\mathcal{P}(^2X)$  is naturally isomorphic t Lip<sub>0</sub>(B<sub>X</sub>), via the restriction map

2

For a Banach space X,  $\mathcal{P}(^2X)$  is a subspace of  $Lip_0(X)$ .

- P(<sup>2</sup>X) is the Banach space of bounded 2-homogeneous polynomials on X.
  - ▶  $P \in \mathcal{P}(^2X)$  if there is a bounded bilinear map  $M: X \times X \to \mathbb{R}$  such that P(x) = M(x, x);
  - $||P||_{\mathcal{P}} = \sup_{x \in B_X} |P(x)|.$
- But polynomials are **not** Lipschitz functions!
- However, they are Lipschitz on the unit ball.
  - Therefore,  $\mathcal{P}(^2X)$  is a natural subspace of *Li*
  - Moreover,  $\|\cdot\|_{\mathcal{P}}$  is equivalent to  $\|\cdot\|_{Lip}$ .
- Consequently, \$\mathcal{P}(^2X)\$ is naturally isomorphic t Lip<sub>0</sub>(B<sub>X</sub>), via the restriction map

2

For a Banach space X,  $\mathcal{P}(^2X)$  is a subspace of  $Lip_0(X)$ .

- P(<sup>2</sup>X) is the Banach space of bounded 2-homogeneous polynomials on X.
  - $P \in \mathcal{P}(^2X)$  if there is a bounded bilinear map  $M: X \times X \to \mathbb{R}$  such that P(x) = M(x, x);
  - $||P||_{\mathcal{P}} = \sup_{x \in B_X} |P(x)|.$
- But polynomials are not Lipschitz functions!
- However, they are Lipschitz on the unit ball.
  - Therefore,  $\mathcal{P}(^2X)$  is a natural subspace of  $Lip_0$
  - Moreover,  $\|\cdot\|_{\mathcal{P}}$  is equivalent to  $\|\cdot\|_{Lip}$ .

 Consequently, P(<sup>2</sup>X) is naturally isomorphic t Lip<sub>0</sub>(B<sub>X</sub>), via the restriction map

2

For a Banach space X,  $\mathcal{P}(^2X)$  is a subspace of  $Lip_0(X)$ .

- P(<sup>2</sup>X) is the Banach space of bounded 2-homogeneous polynomials on X.
  - $P \in \mathcal{P}(^2X)$  if there is a bounded bilinear map  $M: X \times X \to \mathbb{R}$  such that P(x) = M(x, x);
  - $||P||_{\mathcal{P}} = \sup_{x \in B_X} |P(x)|.$
- But polynomials are not Lipschitz functions!
- However, they are Lipschitz on the unit ball.
  - Therefore,  $\mathcal{P}(^2X)$  is a natural subspace of *Lip*
  - Moreover,  $\|\cdot\|_{\mathcal{P}}$  is equivalent to  $\|\cdot\|_{Lip}$

• Consequently,  $\mathcal{P}(^2X)$  is naturally isomorphic to  $Lip_0(B_X)$ , via the restriction map

2

For a Banach space X,  $\mathcal{P}(^2X)$  is a subspace of  $Lip_0(X)$ .

- P(<sup>2</sup>X) is the Banach space of bounded 2-homogeneous polynomials on X.
  - $P \in \mathcal{P}(^2X)$  if there is a bounded bilinear map  $M: X \times X \to \mathbb{R}$  such that P(x) = M(x, x);
  - $||P||_{\mathcal{P}} = \sup_{x \in B_X} |P(x)|.$
- But polynomials are not Lipschitz functions!
- However, they are Lipschitz on the unit ball.
  - Therefore,  $\mathcal{P}(^2X)$  is a natural subspace of  $Lip_0(B_X)$ ;
  - Moreover,  $\|\cdot\|_{\mathcal{P}}$  is equivalent to  $\|\cdot\|_{Lip}$ .

Consequently,  $\mathcal{P}(^2X)$  is naturally isomorphic to  $Lip_0(B_X)$ , via the restriction map

2

For a Banach space X,  $\mathcal{P}(^2X)$  is a subspace of  $Lip_0(X)$ .

- P(<sup>2</sup>X) is the Banach space of bounded 2-homogeneous polynomials on X.
  - ▶  $P \in \mathcal{P}(^2X)$  if there is a bounded bilinear map  $M: X \times X \to \mathbb{R}$  such that P(x) = M(x, x);
  - $||P||_{\mathcal{P}} = \sup_{x \in B_X} |P(x)|.$
- But polynomials are not Lipschitz functions!
- ► However, they are Lipschitz on the unit ball.
  - Therefore,  $\mathcal{P}(^2X)$  is a natural subspace of  $Lip_0(B_X)$ ;
  - Moreover,  $\|\cdot\|_{\mathcal{P}}$  is equivalent to  $\|\cdot\|_{Lip}$ .
- ► Consequently, P(<sup>2</sup>X) is naturally isomorphic to a subspace of Lip<sub>0</sub>(B<sub>X</sub>), via the restriction map

## $P \mapsto P \upharpoonright_{B_X}$ .



The AP passes to complemented subspaces;

- **•** Dineed and Mujica (2015).  $\mathcal{P}(^{2}\ell_{2})$  does not have
- ▶ So, if  $\mathcal{P}({}^{2}\ell_{2}) \subseteq Lip_{0}(B_{\ell_{2}})$  is complemented that have the AP.

#### Question

Is  $\mathcal{P}({}^{2}\ell_{2}) \subseteq Lip_{0}(B_{\ell_{2}})$  a complemented subspace?

- Kaufmann (2015).  $Lip_0(X)$  is isomorphic to
- Thus, a positive answer to this question wou fails to have the AP.



 $B_{\ell_2}$ ) fails

- The AP passes to complemented subspaces;
- ▶ Dineed and Mujica (2015).  $\mathcal{P}(^{2}\ell_{2})$  does not have the AP;
- ▶ So, if  $\mathcal{P}(^{2}\ell_{2}) \subseteq Lip_{0}(B_{\ell_{2}})$  is complemented the have the AP.

#### Question

Is  $\mathcal{P}(^{2}\ell_{2}) \subseteq Lip_{0}(B_{\ell_{2}})$  a complemented subspace?

- Kaufmann (2015).  $Lip_0(X)$  is isomorphic to
- Thus, a positive answer to this question wou fails to have the AP.

- The AP passes to complemented subspaces;
- Dineed and Mujica (2015).  $\mathcal{P}(^{2}\ell_{2})$  does not have the AP;
- So, if P(<sup>2</sup>ℓ<sub>2</sub>) ⊆ Lip<sub>0</sub>(B<sub>ℓ<sub>2</sub></sub>) is complemented, then Lip<sub>0</sub>(B<sub>ℓ<sub>2</sub></sub>) fails to have the AP.

#### Question

Is  $\mathcal{P}(^{2}\ell_{2}) \subseteq Lip_{0}(B_{\ell_{2}})$  a complemented subspace?

- Kaufmann (2015).  $Lip_0(X)$  is isomorphic to Li
- Thus, a positive answer to this question wou fails to have the AP.

- The AP passes to complemented subspaces;
- Dineed and Mujica (2015).  $\mathcal{P}(^{2}\ell_{2})$  does not have the AP;
- So, if P(<sup>2</sup>ℓ<sub>2</sub>) ⊆ Lip<sub>0</sub>(B<sub>ℓ<sub>2</sub></sub>) is complemented, then Lip<sub>0</sub>(B<sub>ℓ<sub>2</sub></sub>) fails to have the AP.

#### Question

Is  $\mathcal{P}({}^{2}\ell_{2}) \subseteq Lip_{0}(\mathcal{B}_{\ell_{2}})$  a complemented subspace?

• Kaufmann (2015).  $Lip_0(X)$  is isomorphic to L

 Thus, a positive answer to this question would fails to have the AP.

- The AP passes to complemented subspaces;
- Dineed and Mujica (2015).  $\mathcal{P}(^{2}\ell_{2})$  does not have the AP;
- So, if P(<sup>2</sup>ℓ<sub>2</sub>) ⊆ Lip<sub>0</sub>(B<sub>ℓ<sub>2</sub></sub>) is complemented, then Lip<sub>0</sub>(B<sub>ℓ<sub>2</sub></sub>) fails to have the AP.

#### Question

Is  $\mathcal{P}(^{2}\ell_{2}) \subseteq Lip_{0}(B_{\ell_{2}})$  a complemented subspace?

• Kaufmann (2015).  $Lip_0(X)$  is isomorphic to  $Lip_0(B_X)$ ;

Thus, a positive answer to this question would fails to have the AP.

- The AP passes to complemented subspaces;
- Dineed and Mujica (2015).  $\mathcal{P}(^{2}\ell_{2})$  does not have the AP;
- So, if P(<sup>2</sup>ℓ<sub>2</sub>) ⊆ Lip<sub>0</sub>(B<sub>ℓ<sub>2</sub></sub>) is complemented, then Lip<sub>0</sub>(B<sub>ℓ<sub>2</sub></sub>) fails to have the AP.

#### Question

Is  $\mathcal{P}(^{2}\ell_{2}) \subseteq Lip_{0}(B_{\ell_{2}})$  a complemented subspace?

- Kaufmann (2015).  $Lip_0(X)$  is isomorphic to  $Lip_0(B_X)$ ;
- ► Thus, a positive answer to this question would yield that  $Lip_0(\ell_2)$  fails to have the AP.

## Question (Repetita iuvant)

Is  $\mathcal{P}(^{2}\ell_{2}) \subseteq Lip_{0}(B_{\ell_{2}})$  a complemented subspace?

#### Theorem (Lindenstrauss, 1964)

 $X^*$  is a 1-complemented subspace of  $Lip_0(X)$ .

• Evidently, 
$$X^* = \mathcal{P}(^1X)$$
;

 The above question is also about the possibilit Lindenstrauss' result to polynomials;

If 'yes', we can answer in the negative Godef

If 'no', Lindenstrauss' result admits no personal version

## Question (Repetita iuvant)

Is  $\mathcal{P}({}^{2}\ell_{2}) \subseteq Lip_{0}(\mathcal{B}_{\ell_{2}})$  a complemented subspace?

#### Theorem (Lindenstrauss, 1964)

 $X^*$  is a 1-complemented subspace of  $Lip_0(X)$ .

• Evidently, 
$$X^* = \mathcal{P}(^1X)$$
;

- The above question is also about the possibility Lindenstrauss' result to polynomials;
- If 'yes', we can answer in the negative Godef
- If 'no', Lindenstrauss' result admits no personal version

## Question (Repetita iuvant)

Is  $\mathcal{P}({}^{2}\ell_{2}) \subseteq Lip_{0}(\mathcal{B}_{\ell_{2}})$  a complemented subspace?

#### Theorem (Lindenstrauss, 1964)

 $X^*$  is a 1-complemented subspace of  $Lip_0(X)$ .

• Evidently, 
$$X^* = \mathcal{P}(^1X)$$
;

 The above question is also about the possibility Lindenstrauss' result to polynomials;

If 'yes', we can answer in the negative Godefront electronic

If 'no', Lindenstrauss' result admits no personal version.

## Question (Repetita iuvant)

Is  $\mathcal{P}({}^{2}\ell_{2}) \subseteq Lip_{0}(\mathcal{B}_{\ell_{2}})$  a complemented subspace?

#### Theorem (Lindenstrauss, 1964)

 $X^*$  is a 1-complemented subspace of  $Lip_0(X)$ .

• Evidently, 
$$X^* = \mathcal{P}(^1X)$$
;

- The above question is also about the possibility to extend Lindenstrauss' result to polynomials;
  - If 'yes', we can answer in the negative Godefrond electric
- If 'no', Lindenstrauss' result admits no petroomal version.

## Question (Repetita iuvant)

Is  $\mathcal{P}({}^{2}\ell_{2}) \subseteq Lip_{0}(\mathcal{B}_{\ell_{2}})$  a complemented subspace?

#### Theorem (Lindenstrauss, 1964)

 $X^*$  is a 1-complemented subspace of  $Lip_0(X)$ .

• Evidently, 
$$X^* = \mathcal{P}(^1X)$$
;

- The above question is also about the possibility to extend Lindenstrauss' result to polynomials;
- If 'yes', we can answer in the negative Godefroy's question;

If 'no', Lindenstrauss' result admits no performant version.

## Question (Repetita iuvant)

Is  $\mathcal{P}({}^{2}\ell_{2}) \subseteq Lip_{0}(\mathcal{B}_{\ell_{2}})$  a complemented subspace?

#### Theorem (Lindenstrauss, 1964)

 $X^*$  is a 1-complemented subspace of  $Lip_0(X)$ .

• Evidently, 
$$X^* = \mathcal{P}(^1X)$$
;

- The above question is also about the possibility to extend Lindenstrauss' result to polynomials;
- If 'yes', we can answer in the negative Godefroy's question;
- If 'no', Lindenstrauss' result admits no polynomial version.



**NO.**  $\mathcal{P}(^{2}\ell_{2}) \subseteq Lip_{0}(B_{\ell_{2}})$  is not complemented.

- We didn't solve the problem we started with
- But, at least, we can tell that this is not the
- Aron and Schottenloher (1976).  $\mathcal{P}({}^{n}\ell_{1})$  is

The result follows from a finite-dimensional, quanti

#### Lheorem (Hájek and R.

Let  $E_n$  be  $\mathbb{R}^n$  with euclidean norm. If Q is any projection from  $Lip_0(B_{E_n})$  onto  $\mathcal{P}(^2E_n)$ , then

$$\|Q\| \ge c \cdot \left(n - 2\sqrt{2}\right)^{1/5}$$



#### **NO.** $\mathcal{P}(^{2}\ell_{2}) \subseteq Lip_{0}(B_{\ell_{2}})$ is not complemented.

- We didn't solve the problem we started with
- But, at least, we can tell that this is not the
- Aron and Schottenloher (1976).  $\mathcal{P}({}^{n}\ell_{1})$  is i

The result follows from a finite-dimensional, quanti

#### Theorem (Hájek and R.

Let  $E_n$  be  $\mathbb{R}^n$  with euclidean norm. If Q is any projection from  $Lip_0(B_{E_n})$  onto  $\mathcal{P}({}^2E_n)$ , then

roach

$$\|Q\| \ge c \cdot \left(n - 2\sqrt{2}\right)^{1/5}$$



**NO.**  $\mathcal{P}(^{2}\ell_{2}) \subseteq Lip_{0}(B_{\ell_{2}})$  is not complemented.

- We didn't solve the problem we started with;
- But, at least, we can tell that this is not the open and the second s
- Aron and Schottenloher (1976).  $\mathcal{P}({}^{n}\ell_{1})$  is is

The result follows from a finite-dimensional, quanti

#### Theorem (Hájek and R.

Let  $E_n$  be  $\mathbb{R}^n$  with euclidean norm. If Q is any projection from  $Lip_0(B_{E_n})$  onto  $\mathcal{P}(^2E_n)$ , then

approach phic to *l* 

$$\|Q\| \ge c \cdot \left(n - 2\sqrt{2}\right)^{1/5}$$



**NO.**  $\mathcal{P}(^{2}\ell_{2}) \subseteq Lip_{0}(B_{\ell_{2}})$  is not complemented.

- We didn't solve the problem we started with;
- But, at least, we can tell that this is not the correct approach.

nic to la

• Aron and Schottenloher (1976).  $\mathcal{P}({}^{n}\ell_{1})$  is is

The result follows from a finite-dimensional, quantitative counter

#### Theorem (Hájek and R.)

Let  $E_n$  be  $\mathbb{R}^n$  with euclidean norm. If Q is any projection from  $Lip_0(B_{E_n})$  onto  $\mathcal{P}({}^2E_n)$ , then

$$\|Q\| \ge c \cdot \left(n - 2\sqrt{2}\right)^{1/5}.$$



**NO.**  $\mathcal{P}(^{2}\ell_{2}) \subseteq Lip_{0}(B_{\ell_{2}})$  is not complemented.

- We didn't solve the problem we started with;
- But, at least, we can tell that this is not the correct approach.
- Aron and Schottenloher (1976).  $\mathcal{P}({}^{n}\ell_{1})$  is isomorphic to  $\ell_{\infty}$ .

The result follows from a finite-dimensional, quantitative counter

#### Theorem (Hájek and R.)

Let  $E_n$  be  $\mathbb{R}^n$  with euclidean norm. If Q is any projection from  $Lip_0(B_{E_n})$  onto  $\mathcal{P}({}^2E_n)$ , then

$$\|Q\| \ge c \cdot \left(n - 2\sqrt{2}\right)^{1/5}.$$



**NO.**  $\mathcal{P}(^{2}\ell_{2}) \subseteq Lip_{0}(B_{\ell_{2}})$  is not complemented.

- We didn't solve the problem we started with;
- But, at least, we can tell that this is not the correct approach.
- Aron and Schottenloher (1976).  $\mathcal{P}({}^{n}\ell_{1})$  is isomorphic to  $\ell_{\infty}$ .

The result follows from a finite-dimensional, quantitative counterpart.

#### Theorem (Hájek and R.)

Let  $E_n$  be  $\mathbb{R}^n$  with euclidean norm. If Q is any projection from  $Lip_0(B_{E_n})$  onto  $\mathcal{P}({}^2E_n)$ , then

$$\|Q\| \ge c \cdot \left(n - 2\sqrt{2}\right)^{1/5}$$

▶ If a Banach space X contains  $(\ell_2^n)_{n=1}^{\infty}$  uniformly complemented, then  $\mathcal{P}(^2X)$  is not complemented in  $Lip_0(B_X)$ ;

▶ **Tzafriri (1974).** If a Banach space admits an unconditional basis, then there is  $p \in \{1, 2, \infty\}$  such that  $(t_{n=1}^{\infty})$  uniformly complemented in X;

If, additionally, X has non-trivial type, it must

#### Corollary/Theorem (Hájek and R.)

If a Banach space X an unconditional basis and non-trivial type, then  $\mathcal{P}(^2X)$  is not complemented in  $Lip_0(B_X)$ .

- $\ell_p \ (1$
- $\blacktriangleright L_p (1$

Recall:  $\mathcal{P}(^{2}\ell_{1})$  is complemented in *Lin* ( $B_{\ell_{1}}$ ) (Aron–Schottenloher).

- ▶ If a Banach space X contains  $(\ell_2^n)_{n=1}^{\infty}$  uniformly complemented, then  $\mathcal{P}(^2X)$  is not complemented in  $Lip_0(B_X)$ ;
- ► Tzafriri (1974). If a Banach space admits an unconditional basis, then there is p ∈ {1, 2, ∞} such that (ℓ<sup>n</sup><sub>p</sub>)<sup>∞</sup><sub>n=1</sub> is uniformly complemented in X;

If, additionally, X has non-trivial type, it must

#### Corollary/Theorem (Hájek and R.)

If a Banach space X an unconditional basis and non-trivial type, then  $\mathcal{P}(^2X)$  is not complemented in  $Lip_0(B_X)$ .

- $\ell_p (1$
- $L_p$  (1

Recall:  $\mathcal{P}(^{2}\ell_{1})$  is complemented in  $Lin(B_{\ell_{1}})$  (Aron–Schottenloher).

- ▶ If a Banach space X contains  $(\ell_2^n)_{n=1}^{\infty}$  uniformly complemented, then  $\mathcal{P}(^2X)$  is not complemented in  $Lip_0(B_X)$ ;
- ► Tzafriri (1974). If a Banach space admits an unconditional basis, then there is p ∈ {1, 2, ∞} such that (ℓ<sup>n</sup><sub>p</sub>)<sup>∞</sup><sub>n=1</sub> is uniformly complemented in X;
- If, additionally, X has non-trivial type, it must be p = 2.

#### Corollary/Theorem (Hájek and R.)

If a Banach space X an unconditional basis and non-trivial type, then  $\mathcal{P}(^2X)$  is not complemented in  $Lip_0(B_X)$ .

- $\ell_p \ (1$
- ▶  $L_p$  (1 <  $p < \infty$ );
- Recall:  $\mathcal{P}(^{2}\ell_{1})$  is complemented in  $Lin_{0}(B_{\ell_{1}})$  (Aron–Schottenloher).

- ▶ If a Banach space X contains  $(\ell_2^n)_{n=1}^{\infty}$  uniformly complemented, then  $\mathcal{P}(^2X)$  is not complemented in  $Lip_0(B_X)$ ;
- ► Tzafriri (1974). If a Banach space admits an unconditional basis, then there is p ∈ {1, 2, ∞} such that (ℓ<sup>n</sup><sub>p</sub>)<sup>∞</sup><sub>n=1</sub> is uniformly complemented in X;
- If, additionally, X has non-trivial type, it must be p = 2.

## Corollary/Theorem (Hájek and R.)

If a Banach space X an unconditional basis and non-trivial type, then  $\mathcal{P}(^2X)$  is not complemented in  $Lip_0(B_X)$ .

- $\ell_p \ (1$
- ▶  $L_p$  (1
- ▶ Recall:  $\mathcal{P}(^{2}\ell_{1})$  is complemented in  $Lip_{0}(B_{\ell_{1}})$  (Aron–Schottenloher).

- ▶ If a Banach space X contains  $(\ell_2^n)_{n=1}^{\infty}$  uniformly complemented, then  $\mathcal{P}(^2X)$  is not complemented in  $Lip_0(B_X)$ ;
- ► Tzafriri (1974). If a Banach space admits an unconditional basis, then there is p ∈ {1, 2, ∞} such that (ℓ<sup>n</sup><sub>p</sub>)<sup>∞</sup><sub>n=1</sub> is uniformly complemented in X;
- If, additionally, X has non-trivial type, it must be p = 2.

## Corollary/Theorem (Hájek and R.)

If a Banach space X an unconditional basis and non-trivial type, then  $\mathcal{P}(^2X)$  is not complemented in  $Lip_0(B_X)$ .

- ▶  $l_p$  (1 < p < ∞);
- $L_p (1$

▶ Recall:  $\mathcal{P}(^{2}\ell_{1})$  is complemented in  $Lip_{0}(B_{\ell_{1}})$  (Aron–Schottenloher).

- ▶ If a Banach space X contains  $(\ell_2^n)_{n=1}^{\infty}$  uniformly complemented, then  $\mathcal{P}(^2X)$  is not complemented in  $Lip_0(B_X)$ ;
- ► Tzafriri (1974). If a Banach space admits an unconditional basis, then there is p ∈ {1, 2, ∞} such that (ℓ<sup>n</sup><sub>p</sub>)<sup>∞</sup><sub>n=1</sub> is uniformly complemented in X;
- If, additionally, X has non-trivial type, it must be p = 2.

#### Corollary/Theorem (Hájek and R.)

If a Banach space X an unconditional basis and non-trivial type, then  $\mathcal{P}(^2X)$  is not complemented in  $Lip_0(B_X)$ .

- ▶  $l_p$  (1 < p < ∞);
- ▶  $L_p$  (1 < p < ∞);

▶ Recall:  $\mathcal{P}(^{2}\ell_{1})$  is complemented in  $Lip_{0}(B_{\ell_{1}})$  (Aron–Schottenloher).

Thank you for your attention !

## The abstract



## The actual talk

