Invent. math. 132, 261-305 (1998)



Higher Bott-Chern forms and Beilinson's regulator

Jose Ignacio Burgos^{1,*}, Steve Wang²

 ¹ Departament D'Álgebra i Geometria. Universitat de Barcelona, Gran Via 585, E-08007 Barcelona, Spain (e-mail burgos@cerber.mat.ub.es)
 ² 700 E. Henry Clay, Apt. 7 Milwaukee, WI 53217, USA

Oblatum 21-III-1997 & 12-VI-1997

Abstract. In this paper, we prove a Gauss-Bonnet theorem for the higher algebraic *K*-theory of smooth complex algebraic varieties. To each exact *n*-cube of hermitian vector bundles, we associate a higher Bott-Chen form, generalizing the Bott-Chern forms associated to exact sequences. These forms allow us to define characteristic classes from *K*-theory to absolute Hodge cohomology. Then we prove that these characteristic classes agree with Beilinson's regulator map.

Introduction

The aim of this paper is to generalize the Gauss-Bonnet theorem to the higher algebraic *K*-theory of smooth complex algebraic varieties.

The Gauss-Bonnet theorem states that, if X is a compact oriented 2-dimensional riemannian manifold, then

$$\iint_X \mathscr{K} dA = 2\pi \chi(X) \;\;,$$

where \mathscr{K} is the gaussian curvature of the riemannian metric and $\chi(X)$ is the Euler characteristic of the manifold X. Thus, this theorem relates a global topological invariant, the Euler characteristic, with a locally defined differential geometrical object, the gaussian curvature. The Gauss-Bonnet theorem may be restated saying that the closed form $\mathscr{K}/2\pi$ represents, in de Rham cohomology, the Euler class of the tangent bundle.

This result was generalized by Chern ([Ch], see also [Gr], [M-S]). If X is an almost complex manifold and E is a complex vector bundle, by topolo-

^{*} Partially supported by the DGICYT nº. PB93-0790

gical means (for instance obstruction theory), we can define some characteristic classes of the vector bundle, called the Chern classes of E,

$$c_i(E) \in H^{2j}(X, (2\pi i)^j \mathbb{Z}), \qquad j \ge 0$$

In this paper we shall restrict the discussion to the Chern character (see [M-S]), denoted $ch_0(E)$, which is a certain power series, with rational coefficients, in the Chern classes. The Chern character is additive for exact sequences. Nevertheless, since the cohomology groups we shall consider will be real vector spaces, any result about the Chern character will imply the analogous result for any power series in the Chern classes.

Let us provide *E* with a hermitian metric *h*. Let E_X denote the differential graded commutative algebra of complex valued differential forms on *X*, and let $E_{X,\mathbb{R}}$ denote the subalgebra of real forms. Let *D* be the unique connection of *E* satisfying

(1) D preserves h.

(2) If $U \subset X$ is an open subset and *s* is a holomorphic section of $E|_U$, then *Ds* is of pure type (1,0).

Let $K = D^2$ be the curvature form. Following Chern and Weil, let us write

(1)
$$\widetilde{\mathrm{ch}}_0(E,h) = \mathrm{tr} \exp(-K) \in \bigoplus_p (2\pi i)^p E_{X,\mathbb{R}}^{p,p}$$

Then the form $\widetilde{ch}_0(E, h)$ is closed. The Chern theorem states that the form $\widetilde{ch}_0(E, h)$ represents, in de Rham cohomology, the Chern character class.

Observe that an oriented riemannian 2 dimensional manifold admits a canonical structure of complex manifold. Since the Euler class is the top Chern class, the Gauss-Bonnet theorem is a particular case of Chern's theorem.

The additivity of the Chern character implies that it induces a group homomorphism

$$\operatorname{ch}_0: K_0(X) \longrightarrow \bigoplus_j (2\pi i)^j H^{2j}(X, \mathbf{Q})$$

where $K_0(X)$ is the Grothendiek group of X.

Characteristic classes for higher algebraic *K*-theory were introduced by Gillet in [Gi]. These classes are defined on any cohomology theory satisfying certain properties, such as de Rham cohomology. Nevertheless, in this case, these higher characteristic classes do not give much information. For instance, for a proper smooth complex algebraic variety, the only non zero classes are the original Chern classes from the K_0 group. In contrast, if the cohomology theory is absolute Hodge cohomology, the Chern character map obtained in this way agrees with the Beilinson regulator map, which is highly non trivial and is involved in very deep and far reaching conjectures [Be].

Recall that, for X a smooth proper algebraic complex variety, we have

$$\begin{aligned} H^{2p}_{\mathscr{H}}(X, \mathbb{R}(p)) &= H^{p,p}(X, \mathbb{C}) \cap (2\pi i)^p H^{2p}(X, \mathbb{R}), \\ H^{2p-1}_{\mathscr{H}}(X, \mathbb{R}(p)) &= H^{p-1,p-1}(X, \mathbb{C}) \cap (2\pi i)^{p-1} H^{2p-2}(X, \mathbb{R}) \end{aligned}$$

Therefore, since, by (1), the Chern character form has the right Hodge type, the Chern theorem implies that the Chern character form represents, in absolute Hodge cohomology, the Chern character class. The question of generalizing the Chern theorem to the characteristic classes from higher K-theory to absolute Hodge cohomology arises naturally.

In their paper about arithmetic characteristic classes of hermitian vector bundles [G-S 1], Gillet and Soulé accomplished the first step of this program, extending the Chern theorem to the case of $K_1(X)$. Let us briefly explain this result.

The elements of $K_1(X)$ may be represented by exact sequences of vector bundles. Thus, the first step is to understand what is the analogue of the Chern forms of hermitian vector bundles in the case of exact sequences of hermitian vector bundles.

Let

$$\overline{\xi}: 0 \longrightarrow (E', h') \longrightarrow (E, h) \longrightarrow (E'', h'') \longrightarrow 0 \quad :$$

be an exact sequence of hermitian vector bundles. Then the Chern character classes satisfy

$$ch_0(E) = ch_0(E') + ch_0(E'')$$
.

Nevertheless, in general, the Chern character form does not behave additively:

$$\widetilde{\mathrm{ch}}_0(E,h) \neq \widetilde{\mathrm{ch}}_0(E',h') + \widetilde{\mathrm{ch}}_0(E'',h'')$$
.

In the case when h' and h'' are the metrics induced by h, Bott and Chern ([B-C]) have defined a differential form, $\widetilde{ch}_1(\overline{\xi})$, which will be called the Bott-Chern form of $\overline{\xi}$, such that

(2)
$$-2\partial\overline{\partial}\widetilde{ch}_{1}(\overline{\xi}) = \widetilde{ch}_{0}(E',h') + \widetilde{ch}_{0}(E'',h'') - \widetilde{ch}_{0}(E,h) .$$

Note that the normalization factor we use is different from the normalization factor used in the original paper. The forms $\widetilde{ch}_1(\overline{\xi})$ are natural and well defined only up to Im ∂ + Im $\overline{\partial}$.

Bismut, Gillet and Soulé [B-G-S], [G-S 1] have given a different construction of Bott-Chern forms that can be applied to the case when h' and h''are not the induced metrics. These Bott-Chern forms are also well defined only up to Im ∂ + Im $\overline{\partial}$. Bott-Chern forms are exactly what we were looking for, and, when X is proper, Gillet and Soulé [G-S 1] have given an explicit description of Beilinson's regulator map for $K_1(X)$ in terms of these Bott-Chern forms, thus extending the Chern theorem to the group K_1 .

As we have seen, the Chern character class on the K_0 is additive for exact sequences. Nevertheless one cannot make a consistent choice of representatives of the Chern character that behave additively for exact sequences. The Bott-Chern forms measure precisely this lack of additivity at the level of Chern forms, and they are responsible for the Chern character for the K_1 -group. Following Schechtman's ideas [Sch] the lack of additivity of the representatives of the Chern character for K_i is responsible for the Chern character for K_{i+1} . Thus, the lack of additivity of Bott-Chern forms should allow us to define second order Bott-Chern forms that give a description of Beilinson's regulator map for the K_2 . And we can repeat this process to obtain Beilinson's regulator map for all the K groups.

In this direction, when X is proper, the second author [Wan] has defined higher Bott-Chern forms for exact hermitian *n*-cubes. These forms may be thought of as an iteration of Bott-Chern forms. Moreover, he has used them to define characteristic classes for higher *K*-theory, proving that, if one can naturally extend higher Bott-Chern forms to the non proper case, then these characteristic classes agree with Beilinson's regulator map.

In this paper we shall give a variant of Wang's original construction that can be easily extended to the non-proper case, thus completing the proof of the Gauss-Bonnet theorem for higher algebraic *K*-theory. An interesting feature of the construction given here is that we obtain well defined Bott-Chern forms and not only modulo Im $\partial + \text{Im }\overline{\partial}$.

Parallel results in the framework of multiplicative *K*-theory have been obtained by Karoubi in [K1] and [K2].

Gillet and Soulé [G-S 1] have used Bott-Chern forms to define arithmetic K_0 groups. Soulé has suggested [So], see also [De], that one may define higher arithmetic *K*-groups as the homotopy fibre of Beilinson's regulator map. We expect that higher Bott-Chern forms, as presented in this paper, will be useful in giving a more concrete definition of higher arithmetic *K*-theory and studying its properties.

Throughout the paper all vector bundles will be algebraic and we shall use the equivalent notion of locally free sheaf.

The plan of the paper is as follows. In Sect. 1 we recall the definition of real absolute Hodge cohomology. We shall also show that real absolute Hodge cohomology can be computed by means of a complex composed by forms defined on $X \times (\mathbb{P}^1)^n$, $n \ge 0$. Higher Bott-Chern forms will live in this complex.

In Sect. 2 we introduce and study some properties of smooth at infinity hermitian metrics. Over a non proper smooth complex variety, to compute real absolute Hodge cohomology, one needs to impose logarithmic conditions at infinity to the differential forms. Thus we cannot use arbitrary hermitian metrics because they will produce differential forms with arbitrary singularities at infinity. The use of smooth at infinity hermitian metrics ensures that Bott-Chern forms have the right behaviour at infinity.

In Sect. 3 we recall the notion of exact metrized *n*-cubes. To each exact *n*-cube, *E*, we shall attach a vector bundle, $\operatorname{tr}_n(E)$ over $X \times (\mathbb{P}^1)^n$ which may be thought of as a homotopy between the faces of *E*. When the hermitian metrics of *E* satisfy certain technical condition, we shall define a natural metric on $\operatorname{tr}_n(E)$. The Chern character form of the vector bundle $\operatorname{tr}_n(E)$ will play the role of higher Bott-Chern forms. Note that these forms live in $X \times (\mathbb{P}^1)^n$.

In Sect. 4 we use higher Bott-Chern forms to define Chern character classes from higher *K*-theory to real absolute Hodge cohomology.

In Sect. 5 we prove that the higher Chern character defined in Sect. 4 agrees with Beilinson's regulator map.

In Sect. 6 we recall several complexes that compute real absolute Hodge cohomology and homology. Using them we give, for X proper, two different versions of higher Bott-Chern forms which are defined on X. The first one, obtained using the Thom-Whitney simple, is multiplicative. The second one agrees with classical Bott-Chern forms and with the original definition due to Wang.

Acknowledgement. We would like to thank Prof. C. Soulé who suggested this question to us and helped us with encouragement and numerous hints. Without him this paper would never have been produced. We would like to thank Prof. V. Navarro Aznar for his help and ideas; in particular, the final definition of Bott-Chern forms is due to a conversation with him. Moreover, he proposed some shortcuts in Sect. 2. We would also like to thank Prof. B. Mazur for his support and guidance. We acknowledge the help of many colleagues for useful conversations which have helped us to understand a number of aspects of the subject. Our thanks to J.B. Bost, N. Dan, H. Gillet, D. Grayson, P. Guillen, F. Lecomte, C. Naranjo, P. Pascual and D. Roessler.

1 Absolute Hodge cohomology

In this section we shall recall the definition of real absolute Hodge cohomology [Be] of a smooth complex algebraic variety X. By a smooth complex variety we shall mean a smooth separated scheme of finite type over \mathbb{C} . We shall also construct a complex, composed by forms on $X \times (\mathbb{P}^1)^n$, $n \ge 0$, whose cohomology is the real absolute Hodge cohomology of X.

(1.1) Let \overline{X} be a smooth proper complex variety. Let $Y \subset \overline{X}$ be a normal crossing divisor and let us write $X = \overline{X} - Y$. Let $E_{\overline{X}}^*$ be the differential graded algebra of differential forms on \overline{X} , and let $E_{\overline{X}}^*(\log Y)$ be the differential graded algebra of \mathbb{C}^∞ complex differential forms on \overline{X} with logarithmic singularities along Y (see [Bu 1]). The algebra $E_{\overline{X}}^*(\log Y)$ has a real structure, $E_{\overline{X}}^*(\log Y)_{\mathbb{R}}$, a weigh filtration W defined over \mathbb{R} and a Hodge filtration F. Moreover the cohomology of this algebra gives us the cohomology of X with its real mixed Hodge structure.

Let us denote by \widehat{W} the décalée filtration of W. That is

$$\widehat{W}_{r}E_{\overline{X}}^{n}(\log Y) = \left\{ x \in W_{r-n}E_{\overline{X}}^{n}(\log Y) \mid dx \in W_{r-n-1}E_{\overline{X}}^{n+1}(\log Y) \right\}$$

We write

$$E^*_{\log}(X) = \lim_{\stackrel{\longrightarrow}{(\tilde{X}_{\alpha}, Y_{\alpha})}} E^*_{\tilde{X}_{\alpha}}(\log Y_{\alpha}) ,$$

where the limit is taken along all the smooth compactifications \widetilde{X}_{α} of X with $Y_{\alpha} = \widetilde{X}_{\alpha} - X$ a normal crossing divisor. Then $E^*_{\log}(X)$ is a differential graded algebra and it has an induced real structure, a weight filtration and a Hodge filtration. Moreover the map

$$\left(E^*_{\overline{X}}(\log Y)_{\mathbb{R}}, \widehat{W}\right) \longrightarrow \left(E^*_{\log}(X)_{\mathbb{R}}, \widehat{W}\right)$$

is a filtered quasi-isomorphism and the map

$$\left(E^*_{\overline{X}}(\log Y), \widehat{W}, F\right) \longrightarrow \left(E^*_{\log}(X), \widehat{W}, F\right)$$

is a bifiltered quasi-isomorphism.

(1.2) Let us write

$$\mathfrak{H}^*(X,p) = s((2\pi i)^p \widehat{W}_{2p} E^*_{\log}(X)_{\mathbb{R}} \oplus \widehat{W}_{2p} \cap F^p E^*_{\log}(X) \xrightarrow{u} \widehat{W}_{2p} E^*_{\log}(X)) ,$$

where u(r, f) = f - r and s denotes the simple of a morphism of complexes, i.e. the cône shifted by one. The differential of this complex will be denoted by $d_{\mathfrak{H}}$.

The real absolute Hodge cohomology of X [Be] is

$$H^n_{\mathscr{H}}(X, \mathbb{R}(p)) = H^n(\mathfrak{H}(X, p))$$

(1.3) A cubical or cocubical object (see [G-N-P-P]) is an object modeled on the cube in the same way as a simplicial or cosimplicial object is modeled on the simplex. Let $(\mathbb{P}^1_{\mathbb{C}})^{\cdot}$ be the cocubical scheme which in degree *n* is $(\mathbb{P}^1_{\mathbb{C}})^n$, the *n*-fold product of the complex projective line. The faces and degeneracies

$$\begin{aligned} d_j^i : \left(\mathbf{P}_{\mathbb{C}}^1 \right)^n &\longrightarrow \left(\mathbf{P}_{\mathbb{C}}^1 \right)^{n+1}, \quad i = 1, \dots, n+1, \ j = 0, 1\\ s^i : \left(\mathbf{P}_{\mathbb{C}}^1 \right)^n &\longrightarrow \left(\mathbf{P}_{\mathbb{C}}^1 \right)^{n-1}, \quad i = 1, \dots, n \end{aligned}$$

are given by

$$d_0^i(x_1, \dots, x_n) = (x_1, \dots, x_{i-1}, (0:1), x_i, \dots, x_n)$$

$$d_1^i(x_1, \dots, x_n) = (x_1, \dots, x_{i-1}, (1:0), x_i, \dots, x_n)$$

$$s^i(x_1, \dots, x_n) = (x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)$$

266

(1.4) The complexes $\mathfrak{H}^*(X \times (\mathbb{P}^1)^{\cdot}, p)$ form a cubical complex. We shall write

$$s_i = \left(\mathrm{Id} \times s^i
ight)^* ,$$

 $d_i^j = \left(\mathrm{Id} \times d_j^i
ight)^* .$

Let us denote by $\mathfrak{H}^{*,*}_{\mathbb{P}}(X,p)$ the associated double complex. That is

$$\mathfrak{H}^{r,n}_{\mathbb{P}}(X,p) = \mathfrak{H}^{r}(X \times (\mathbb{P}^{1})^{-n},p) ,$$

with differentials

$$d' = d_{\mathfrak{H}},$$

 $d'' = \sum (-1)^{i+j} d_i^j.$

(1.5) We want to obtain from $\mathfrak{H}_{\mathbb{P}}^{*,*}(X,p)$, a complex which computes the absolute Hodge cohomology of X. On the one hand, since we are using a cubical theory we need to factor out by the degenerate elements (see [Mas]). On the other hand, we need to kill all cohomology classes coming from the projective spaces.

Let us denote by $p_0: X \times (\mathbb{P}^1)^n \longrightarrow X$ the projection over the first factor and by $p_i: X \times (\mathbb{P}^1)^n \longrightarrow \mathbb{P}^1$, i = 1, ..., n, the projection over the *i*-th projective line.

Let (x : y) be homogeneous coordinates of \mathbb{R}^1 . Let us write

$$g = \log \frac{x\bar{x} + y\bar{y}}{x\bar{x}}$$
 .

Let $\omega = \partial \overline{\partial} g \in (2\pi i) E^2_{\mathbb{P}^1,\mathbb{R}}$ be a Kähler form over \mathbb{P}^1 . Let $\omega_i = p_i^* \omega \in E^*_{\log}(X \times (\mathbb{P}^1)^n)$. For an element

$$x = (r, f, \eta) \in \mathfrak{H}^r (X \times (\mathbb{P}^1)^n, p)$$
,

we shall write

$$\omega_i \wedge x = (\omega_i \wedge r, \omega_i \wedge f, \omega_i \wedge \eta)$$

$$\in \mathfrak{H}^{r+2} (X \times (\mathbb{P}^1)^n, p+1) .$$

Definition 1.1. We shall denote by $\tilde{\mathfrak{H}}^{*,*}(X,p)$ the double complex given by

$$\widetilde{\mathfrak{H}}^{r,n}(X,p) = \mathfrak{H}^{r,n}(X,p) \bigg/ \sum_{i=1}^{n} s_i \Big(\mathfrak{H}^{r,n+1}(X,p) \Big) \oplus \omega_i \wedge s_i \Big(\mathfrak{H}^{r-2,n+1}(X,p-1) \Big) \quad .$$

We shall denote by $\tilde{\mathfrak{H}}^*(X,p)$ the associated simple complex. The differential of this complex will be denoted by d.

In the definition of $\tilde{\mathfrak{F}}^{r,n}(X,p)$, the first summand of the quotient is meant to kill the degenerate classes, whereas the second summand should kill the classes coming from the projective spaces. The next result shows that we have reached our objective.

Proposition 1.2. The natural morphism of complexes

$$\iota:\mathfrak{H}^*(X,p)=\widetilde{\mathfrak{H}}^{*,0}(X,p)\longrightarrow \widetilde{\mathfrak{H}}^*(X,p)$$

is a quasi-isomorphism.

Proof. Since $\tilde{\mathfrak{F}}^*(X,p)$ is a simple complex associated to a double complex, there is a second quadrant spectral sequence with E_1 term

$$E_1^{r,n} = H^r\Big(\widetilde{\mathfrak{H}}^{*,n}(X,p)\Big)$$
 .

When this spectral sequence converges, the limit is $H^*(\tilde{\mathfrak{H}}^*(X,p))$. The following lemma shows that this spectral sequence converges and implies that *i* is a quasi-isomorphism.

Lemma 1.3. For n < 0 the cohomology of the complex $\tilde{\mathfrak{H}}^{*,n}(X,p)$ is zero.

Proof. For each j let us write

$$\widetilde{\mathfrak{H}}_{j}^{r,n}(X,p) = \mathfrak{H}_{\mathbb{P}}^{r,n}(X,p) \bigg/ \sum_{i=1}^{j} s_i \Big(\mathfrak{H}_{\mathbb{P}}^{r,n+1}(X,p) \Big) \oplus \omega_i \wedge s_i \Big(\mathfrak{H}_{\mathbb{P}}^{r-2,n+1}(X,p-1) \Big) .$$

Let us prove, by induction over *j*, that for $j \ge 1$

$$H^*\left(\widetilde{\mathfrak{H}}_j^{*,n}(X,p)\right) = 0$$
.

For j = 1, $n \leq -1$, the complex $\tilde{\mathfrak{H}}_1^{r,n}(X,p)$ is the cokernel of the monomorphism

$$\mathfrak{H}^*(X \times (\mathbb{P}^1)^{-n-1}, p) \oplus \mathfrak{H}^*(X \times (\mathbb{P}^1)^{-n-1}, p-1)[-2] \longrightarrow \mathfrak{H}^*(X \times (\mathbb{P}^1)^{-n}, p)$$

$$\alpha \qquad \oplus \qquad \beta \qquad \longmapsto s_1(\alpha) + \omega_1 \wedge s_1(\beta)$$

But by the Dold-Thom isomorphism for absolute Hodge cohomology, the above morphism is a quasi-isomorphism. For j > 1, n < -1, $\tilde{\mathfrak{G}}_{j}^{*,n}(X,p)$ is the cokernel of the monomorphism

$$\begin{array}{ccc} \widetilde{\mathfrak{H}}_{j-1}^{*,n-1}(X,p) \ \oplus \ \widetilde{\mathfrak{H}}_{j-1}^{*,n-1}(X,p-1)[-2] \ \longrightarrow \ & \widetilde{\mathfrak{H}}_{j-1}^{*,n}(X,p) \\ \alpha \ \oplus \ & \beta \ & \longmapsto \ & s_j(\alpha) + \omega_j \wedge s_j(\beta) \ . \end{array}$$

By induction hypothesis, the source and the target of this morphism have zero cohomology. Therefore the cokernel also has zero cohomology.

2 Smooth at infinity hermitian metrics

In this section we introduce smooth at infinity hermitian metrics. For a smooth complex variety X and a locally free sheaf \mathcal{F} , a smooth at infinity hermitian metric is a metric that can be extended to a smooth metric over some compactification of \mathcal{F} . The interest of smooth at infinity hermitian metrics is that they provide representatives of Chern classes in absolute Hodge cohomology.

(2.1) Before defining smooth at infinity hermitian metrics, we shall study classes of compactifications of locally free sheaves.

Definition 2.1. Let X be a smooth complex variety and let \mathcal{F} be a locally free sheaf over X. A compactification of \mathcal{F} is a smooth compactification of X, $i: X \longrightarrow \widetilde{X}$, a locally free sheaf $\widetilde{\mathscr{F}}$ over \widetilde{X} and an isomorphism $\varphi: \mathscr{F} \longrightarrow i^* \widetilde{\mathscr{F}}$. A compactification of \mathcal{F} will be denoted by $(i, \widetilde{X}, \widetilde{\mathcal{F}}, \varphi)$. Usually we shall identify X with i(X) and \mathscr{F} with $\widetilde{\mathscr{F}}|_X$, and denote a compactification by $(\widetilde{\mathscr{F}},\widetilde{X}).$

Proposition 2.2. Let X be a smooth complex variety and let \mathcal{F} be a locally free sheaf over X. Then there exists a compactification of \mathcal{F} .

Proof. Let $X \longrightarrow \widetilde{X}_1$ be any compactification of X. Then there is a coherent sheaf $\widetilde{\mathscr{F}}_1$ on \widetilde{X}_1 such that $\widetilde{\mathscr{F}}_1|_X = \mathscr{F}$. By [Ro] (see also [Ri] and [N 1]) there is a proper modification $\psi: \widetilde{X} \longrightarrow \widetilde{X}_1$, which induces an isomorphism $\psi^{-1}(X) \longrightarrow X$, and such that $\widetilde{\mathscr{F}} = \psi^*(\widetilde{\mathscr{F}}_1) / \operatorname{Tor}(\psi^*(\widetilde{\mathscr{F}}_1))$ is a locally free sheaf. Moreover $\widetilde{\mathscr{F}}|_{\psi^{-1}(X)}$ is isomorphic to $\widetilde{\mathscr{F}}_1|_X$. Thus the induced map $i: X \longrightarrow \widetilde{X}$ is a compactification of X, and $\widetilde{\mathscr{F}}$ is a compactification of \mathscr{F} .

Definition 2.3. Let X be a smooth complex variety and let \mathcal{F} be a locally free sheaf over X. Let $(i_1, \widetilde{X}_1, \widetilde{\mathscr{F}}_1, \varphi_1)$ and $(i_2, \widetilde{X}_2, \widetilde{\mathscr{F}}_2, \varphi_2)$ be two compactifications of \mathcal{F} . We say that $\widetilde{\mathcal{F}}_1$ and $\widetilde{\mathcal{F}}_2$ are equivalent if there exists a third compactification $(i_3, \widetilde{X}_3, \widetilde{\mathscr{F}}_3, \varphi_3)$ and morphisms $\psi_1 : \widetilde{X}_3 \longrightarrow \widetilde{X}_1$ and $\psi_2 : \widetilde{X}_3 \longrightarrow \widetilde{X}_2$ such that

 ψ₁ ∘ i₃ = i₁ and ψ₂ ∘ i₃ = i₂.
 There are isomorphisms α₁ : ℱ₃ → ψ₁^{*}ℱ₁ and α₂ : ℱ₃ → ψ₂^{*}ℱ₂ such that $i_3^* \alpha_1 \circ \varphi_3 = \varphi_1$ and $i_3^* \alpha_2 \circ \varphi_3 = \varphi_2$.

In order to simplify the notation, a class of equivalent compactifications of \mathscr{F} will be denoted by a single symbol, for instance $\widetilde{\mathscr{F}}$. Moreover, if there is no danger of confusion, we shall denote by the same symbol the locally free sheaf which appears in any representative of this class.

(2.2) Let us see that a compactification class induces uniquely determined compactification classes in quotients and subsheaves.

Theorem 2.4. Let X be a smooth complex variety and let

$$\xi: \ 0 \longrightarrow \mathscr{F} \xrightarrow{f} \mathscr{G} \xrightarrow{g} \mathscr{H} \longrightarrow 0$$

be an exact sequence of locally free sheaves over X. Then for any compactification class $\tilde{\mathcal{G}}$ of \mathcal{G} , there are uniquely determined compactification classes $\tilde{\mathcal{F}}$ and $\tilde{\mathcal{H}}$ of \mathcal{F} and \mathcal{H} respectively such that ξ extends to an exact sequence

$$\widetilde{\xi}: \ 0 \longrightarrow \widetilde{\mathscr{F}} \xrightarrow{\widetilde{f}} \widetilde{\mathscr{G}} \xrightarrow{\widetilde{g}} \widetilde{\mathscr{H}} \longrightarrow 0 \ ,$$

over a compactification \widetilde{X} of X.

Proof. Let \widetilde{X}_1 be a compactification of X where $\widetilde{\mathscr{G}}$ is defined. Let $r = \operatorname{rk} \mathscr{H}$. Let $\operatorname{Grass}_{\widetilde{X}_i}(\widetilde{\mathscr{G}})$ be the Grassmanian of rank r quotients of $\widetilde{\mathscr{G}}$ [G-D]. Let us denote by \mathscr{U} the universal bundle on $\operatorname{Grass}_{\widetilde{X}_i}(\widetilde{\mathscr{G}})$. The exact sequence

$$\xi: \ 0 \longrightarrow \mathscr{F} \xrightarrow{f} \mathscr{G} \xrightarrow{g} \mathscr{H} \longrightarrow 0$$

induces a morphism

$$\varphi: X \longrightarrow \mathbf{Grass}_{\widetilde{X}_1}^r(\widetilde{\mathscr{G}})$$
.

By resolution of singularities, there is a proper modification \widetilde{X} of \widetilde{X}_1 , which is a compactification of X and such that φ extends to a morphism

$$\widetilde{\varphi}: \widetilde{X} \longrightarrow \mathbf{Grass}_{\widetilde{X}_1}(\widetilde{\mathscr{G}})$$
.

Then $\widetilde{\mathscr{H}} = \widetilde{\varphi}^*(\mathscr{U})$ is a compactification of $\mathscr{H}, \ \widetilde{\mathscr{F}} = \operatorname{Ker}(\widetilde{\mathscr{G}} \longrightarrow \widetilde{\mathscr{H}})$ is a compactification of \mathscr{F} and ξ extends to an exact sequence

$$\widetilde{\xi}: \ 0 \longrightarrow \widetilde{\mathscr{F}} \xrightarrow{\widetilde{f}} \widetilde{\mathscr{G}} \xrightarrow{\widetilde{g}} \widetilde{\mathscr{H}} \longrightarrow 0$$

The unicity follows from the fact that, since X is dense in \tilde{X} , the morphism $\tilde{\varphi}$ is unique.

Definition 2.5. Let X be a smooth complex variety and let

$$\xi: \ 0 \longrightarrow \mathscr{F} \xrightarrow{f} \mathscr{G} \xrightarrow{g} \mathscr{H} \longrightarrow 0$$

be an exact sequence of locally free sheaves over X. Let $\widetilde{\mathscr{G}}$ be a class of compactifications of \mathscr{G} . Then the classes of compactifications $\widetilde{\mathscr{F}}$ and $\widetilde{\mathscr{H}}$, of \mathscr{F} and \mathscr{H} respectively, obtained in theorem 2.4 are called the induced compactifications.

(2.3) Let us introduce smooth at infinity hermitian metrics.

Definition 2.6. Let X be a smooth complex variety, let \mathcal{F} be a locally free sheaf over X and let h be an hermitian metric on \mathcal{F} . We say that h is smooth at infinity if there exist a compactification $\tilde{\mathcal{F}}$ of \mathcal{F} , and a smooth metric \tilde{h} on $\tilde{\mathcal{F}}$ such that $\tilde{h}|_X = h$.

A smooth at infinity hermitian metric determines univocally a compactification class.

Proposition 2.7. Let X be a smooth complex variety and let \mathcal{F} be a locally free sheaf on X. Let $\tilde{\mathcal{F}}$ and $\tilde{\mathcal{F}}'$ be two compactifications of \mathcal{F} and let \tilde{h} and \tilde{h}' be smooth metrics on $\tilde{\mathcal{F}}$ and $\tilde{\mathcal{F}}'$. If $\tilde{h}|_X = \tilde{h}'|_X$, then $\tilde{\mathcal{F}}$ and $\tilde{\mathcal{F}}'$ are equivalent compactifications.

Proof. We can assume that both compactifications are defined over the same variety \widetilde{X} . Let $\mathscr{K}_{\widetilde{X}}$ be the sheaf of rational functions over \widetilde{X} .

The identity on \mathcal{F} induces morphisms

$$\begin{split} f: \widetilde{\mathscr{F}} \otimes \mathscr{K}_{\widetilde{X}} & \longrightarrow \widetilde{\mathscr{F}}' \otimes \mathscr{K}_{\widetilde{X}} \ , \\ f': \widetilde{\mathscr{F}}' \otimes \mathscr{K}_{\widetilde{X}} & \longrightarrow \widetilde{\mathscr{F}} \otimes \mathscr{K}_{\widetilde{X}} \ , \end{split}$$

which are inverses of each other. By symmetry it is enough to show that $f(\widetilde{\mathscr{F}}) \subset \widetilde{\mathscr{F}}'$.

Let U be a Zariski open subset of \widetilde{X} . A section $s \in \Gamma(U, \widetilde{\mathscr{F}}' \otimes \mathscr{K}_{\widetilde{X}})$ belongs to $\Gamma(U, \widetilde{\mathscr{F}}')$ if and only if $\widetilde{h}'(s(x)) < \infty$ for all $x \in U$. But if $s \in \Gamma(U, \widetilde{\mathscr{F}})$ then $\widetilde{h}'(f(s))|_{X \cap U} = \widetilde{h}(s)|_{X \cap U}$. Since $U \cap X$ is dense in U we have $\widetilde{h}'(f(s(x))) = \widetilde{h}(s(x)) < \infty$ for all $x \in U$.

Proposition 2.8. Let

$$\xi: 0 \longrightarrow \mathscr{F}' \longrightarrow \mathscr{F} \longrightarrow \mathscr{F}'' \longrightarrow 0$$

be an exact sequence of locally free sheaves on X and let h be a smooth at infinity metric on \mathcal{F} . Then the metrics h' and h'' induced by h in \mathcal{F}' and \mathcal{F}'' are smooth at infinity.

Proof. Let $\widetilde{\mathscr{F}}$ be a compactification of \mathscr{F} provided with a metric \widetilde{h} , such that $\widetilde{h}|_{\chi} = h$. By theorem 2.4. there are compactifications $\widetilde{\mathscr{F}}'$ and $\widetilde{\mathscr{F}}''$ such that ξ can be extended to an exact sequence

$$\widetilde{\xi}: 0 \longrightarrow \widetilde{\mathscr{F}}' \longrightarrow \widetilde{\mathscr{F}} \longrightarrow \widetilde{\mathscr{F}}'' \longrightarrow 0 \quad .$$

Then the metric \tilde{h} induces smooth metrics \tilde{h}' and \tilde{h}'' on $\tilde{\mathscr{F}}'$ and $\tilde{\mathscr{F}}''$. But the restrictions of \tilde{h}' and \tilde{h}'' to X are h' and h''. Therefore these metrics are smooth at infinity.

Proposition 2.9. Let $f: X \longrightarrow Y$ be a morphism between smooth complex varieties. Let (\mathcal{F}, h) be a locally free sheaf over Y with h a smooth at infinity metric. Then (f^*h) is a smooth at infinity metric on the locally free sheaf $f^*\mathcal{F}$.

Proof. Let $(\widetilde{Y}, \widetilde{\mathscr{F}})$ be a compactification of (Y, \mathscr{F}) , such that there exists a hermitian metric \widetilde{h} with $\widetilde{h}|_X = h$. Let \widetilde{X}_1 be any compactification of X. We shall denote by Γ the graph of f, and by $\overline{\Gamma}$ the adherence of Γ in $\widetilde{X}_1 \times \widetilde{Y}$. Let \widetilde{X} be a resolution of singularities of $\overline{\Gamma}$ and let $\widetilde{f} : \widetilde{X} \longrightarrow \widetilde{Y}$ be the induced morphism. Then $(\widetilde{X}, \widetilde{f}^* \widetilde{\mathscr{F}})$ is a compactification of $(X, f^* \mathscr{F})$ and $\widetilde{f}^* \widetilde{h}$ is a smooth metric such that $\widetilde{f}^* \widetilde{h}|_X = f^* h$. Therefore $f^* h$ is smooth at infinity.

(2.4) Let us see that smooth at infinity hermitian metrics provide representatives of the Chern character classes in absolute Hodge cohomology. Let X be a smooth complex variety, \mathscr{F} a locally free sheaf and h a smooth at infinity hermitian metric. Let $\widetilde{\mathscr{F}}$ be the compactification class of \mathscr{F} determined by h, \widetilde{X} a compactification of X where $\widetilde{\mathscr{F}}$ is defined, and \widetilde{h} a smooth metric on $\widetilde{\mathscr{F}}$ extending h. Let K (resp. \widetilde{K}) be the curvature form of (\mathscr{F}, h) (resp. $(\widetilde{\mathscr{F}}, \widetilde{h})$). Let us write

$$\widetilde{\mathrm{ch}}_0(\mathscr{F},h) = \mathrm{Tr} \, \exp(-K) \; ,$$

 $\widetilde{\mathrm{ch}}_0(\widetilde{\mathscr{F}},\widetilde{h}) = \mathrm{Tr} \, \exp(-\widetilde{K}) \; .$

These forms are closed. Moreover,

$$\widetilde{\mathrm{ch}}_0(\widetilde{\mathscr{F}},\widetilde{h})\in \bigoplus (2\pi i)^p E^{p,p}_{\widetilde{X}|\mathbb{R}}$$
 .

Since $\widetilde{ch}_0(\widetilde{\mathscr{F}},\widetilde{h})|_X = \widetilde{ch}_0(\mathscr{F},h),$

$$\widetilde{\mathrm{ch}}_0(\mathscr{F},h) \in \bigoplus_{p \ge 0} \Big(W_0 E_{\mathrm{log}}^{2p}(X)_{\mathbb{R}} \cap F^p E_{\mathrm{log}}^{2p}(X) \Big) \; .$$

Since this form is closed,

$$\widetilde{\mathrm{ch}}_{0}(\mathscr{F},h) \in \bigoplus_{p\geq 0} \left(\widehat{W}_{2p} E^{2p}_{\log}(X)_{\mathbb{R}} \cap F^{p} E^{2p}_{\log}(X) \right) \;.$$

Thus the triple

$$\widetilde{\mathrm{ch}}_0(\mathscr{F},h)_{\mathscr{H}} = (\widetilde{\mathrm{ch}}_0(\mathscr{F},h),\widetilde{\mathrm{ch}}_0(\mathscr{F},h),0)$$

is a cycle of $\bigoplus_{p\geq 0} \mathfrak{H}^{2p}(X,p)$.

Proposition 2.10. The cycle $\widetilde{ch}_0(\mathscr{F}, h)_{\mathscr{H}}$ represents the Chern character of \mathscr{F} in absolute Hodge cohomology.

Proof. If X is proper we have

$$H^{2p}_{\mathscr{H}}(X, \mathbb{R}(p)) = H^{p,p}(X, (2\pi i)^p \mathbb{R})$$

Therefore the result follows from the classical description of the Chern character in terms of curvature forms. In the non proper case it follows from the functoriality of the Chern character.

3 Exact *n*-cubes of locally free sheaves

In this section we shall recall the notion of exact *n*-cube (see [Lo 2], [Wan]). To each metrized exact *n*-cube, $\overline{\mathscr{F}}$, which satisfies certain conditions, we shall associate a metrized locally free sheaf on $X \times (\mathbb{P}^1)^n$, called the *n*-th transgression of $\overline{\mathscr{F}}$. This transgression can be viewed as a homotopy between its vertexes. The Chern character form of the transgression will play the role of higher Bott-Chern forms.

(3.1) First some notations. Let $\langle -1, 0, 1 \rangle$ be the category associated to the ordered set $\{-1, 0, 1\}$. Let $\langle -1, 0, 1 \rangle^n$ be its *n*-th cartesian power. By convention, the category $\langle -1, 0, 1 \rangle^0$ has one element and one morphism.

Let & be an exact category.

Definition 3.1. A *n*-cube of \mathfrak{E} , \mathscr{F} , is a functor from $\langle -1, 0, 1 \rangle^n$ to \mathfrak{E} .

Definition 3.2. Given a n-cube \mathscr{F} , and numbers $i \in \{1, ..., n\}$, $j \in \{-1, 0, 1\}$, then the n - 1-cube, $\partial_i^j \mathscr{F}$ defined by

$$\left(\partial_{i}^{j}\mathscr{F}\right)_{\alpha_{1},\ldots,\alpha_{n-1}}=\mathscr{F}_{\alpha_{1},\ldots,\alpha_{i-1},j,\alpha_{i},\ldots,\alpha_{n-1}}$$

is called an face of \mathscr{F} . Given a number $i \in \{1, \ldots, n\}$ and a n-1-tuple $\alpha = (\alpha_1, \ldots, \alpha_{n-1}) \in \{-1, 0, 1\}^{n-1}$, the sequence

$$\partial_{i^c}^{\alpha}\mathscr{F} = \partial_n^{\alpha_{n-1}} \dots \partial_{i+1}^{\alpha_i} \partial_{i-1}^{\alpha_{i-1}} \dots \partial_1^{\alpha_1} \mathscr{F}$$

is called an edge of \mathcal{F} .

Explicitly, the edge $\partial_{i^c}^{\alpha} \mathscr{F}$ *is*

$$\mathscr{F}_{\alpha_1,\dots,\alpha_{i-1},-1,\alpha_i,\dots,\alpha_{n-1}}\longrightarrow \mathscr{F}_{\alpha_1,\dots,\alpha_{i-1},0,\alpha_i,\dots,\alpha_{n-1}}\longrightarrow \mathscr{F}_{\alpha_1,\dots,\alpha_{i-1},1,\alpha_i,\dots,\alpha_{n-1}}$$

Definition 3.3. A n-cube is called an exact n-cube if all its edges are short exact sequences.

We shall denote by $\underline{C}_n \mathfrak{E}$ the exact category of exact n-cubes. Observe that, for all non negative integers n, m, there is a natural isomorphism of categories $\underline{C}_n \underline{C}_m \mathfrak{E} \longrightarrow \underline{C}_{n+m} \mathfrak{E}$. In particular, an exact n-cube can be viewed as an exact sequence of exact n - 1-cubes or as an exact n - 1-cube of exact sequences. The maps

$$\partial_i^j : \operatorname{Ob} \underline{C}_n \mathfrak{E} \longrightarrow \operatorname{Ob} \underline{C}_{n-1} \mathfrak{E}$$
,

are called face maps. The maps

 s_i^j : Ob $\underline{C}_n \mathfrak{E} \longrightarrow$ Ob $\underline{C}_{n+1} \mathfrak{E}$, for $i = 1, \dots, n$, and j = -1, 1,

given by

$$s_i^j(\mathscr{F})_{\alpha_1,\ldots,\alpha_{n+1}} = \begin{cases} 0, & \text{if } \alpha_i = j, \\ \mathscr{F}_{\alpha_1},\ldots,\alpha_{i-1},\alpha_{i+1},\ldots,\alpha_{n+1}, & \text{if } \alpha_i \neq j \end{cases}$$

are called degeneracy maps. An exact n-cube $\mathscr{F} \in \text{Im } s_i^j$ is called degenerate.

(3.2) We shall write $C_n \mathfrak{E} = \text{Ob } \underline{C}_n \mathfrak{E}$ and $C \mathfrak{E} = \coprod C_n \mathfrak{E}$.

Assume that the category \mathfrak{E} is small. To avoid set theoretical problems, in the sequel we shall always assume tacitly that we replace any large category by an equivalent small full subcategory. Observe that the diagram $C\mathfrak{E}$ behaves like a cubical diagram. We have replaced the category $\langle 0, 1 \rangle$ by the category $\langle -1, 0, 1 \rangle$. This motivates the following construction.

Let $\mathbb{Z}C_n\mathfrak{E}$ be the free abelian group generated by $C_n\mathfrak{E}$. And let the differential $d: \mathbb{Z}C_n\mathfrak{E} \longrightarrow \mathbb{Z}C_{n-1}\mathfrak{E}$ be given by

$$d = \sum_{i=1}^{n} \sum_{j=-1}^{1} (-1)^{i+j} \partial_i^j$$
.

Let $D_n \subset \mathbb{Z}C_n \mathfrak{E}$ be the subgroup generated by the degenerate exact *n*-cubes. Then $dD_n \subset D_{n-1}$. Therefore the following definition makes sense.

Definition 3.4. The homology complex associated to $C\mathfrak{E}$ is

$$\mathbb{Z}C\mathfrak{E} = \mathbb{Z}C\mathfrak{E}/D$$
.

(3.3) For the remainder of the section, let us fix a smooth complex variety X. Let $\mathfrak{E}(X)$ be the exact category of locally free sheaves on X and let $\overline{\mathfrak{E}}(X)$ be the exact category of pairs (\mathcal{F}, h) , where $\mathcal{F} \in \operatorname{Ob} \mathfrak{E}(X)$ and h is a smooth at infinity hermitian metric on \mathcal{F} . The morphisms of this category are

$$\operatorname{Hom}_{\overline{\mathfrak{E}}(X)}((\mathscr{F},h),(\mathscr{F}',h')) = \operatorname{Hom}_{\mathfrak{E}(X)}(\mathscr{F},\mathscr{F}')$$

Let \mathscr{F} be the forgetful functor $\overline{\mathfrak{G}}(X) \longrightarrow \mathfrak{G}(X)$. By choosing metrics we may construct a functor $G : \mathfrak{G}(X) \longrightarrow \overline{\mathfrak{G}}(X)$. Then $F \circ G$ is the identify functor on the category $\mathfrak{G}(X)$. Moreover, the identity morphism on the vector bundles

is a natural transformation between $G \circ F$ and the identity functor in $\overline{\mathfrak{E}}(X)$. Thus *F* is an equivalence of categories.

For simplicity we shall write $C(X) = C\overline{\mathfrak{E}}(X)$. An element $\overline{\mathscr{F}} \in C_n(X)$ is called a metrized exact *n*-cube of locally free sheaves.

(3.4) For technical reasons we need to work with metrized exact *n*-cubes which have, in all the quotients, the induced metrics.

Definition 3.5. We shall say that a metrized exact n-cube, $\overline{\mathscr{F}} = \{(\mathscr{F}_{\alpha}, h_{\alpha})\}$ has induced quotient metrics (an emi-n-cube for short) if, for each n-tuple $\alpha = (\alpha_1, \ldots, \alpha_n)$, and each i with $\alpha_i = 1$, the metric h_{α} is induced by the metric $h_{(\alpha_1,\ldots,\alpha_{i-1},0,\alpha_{i+1},\ldots,\alpha_n)}$.

Let us see that there are enough emi-*n*-cubes. Let $\alpha \in \{-1, 0, 1\}^n$ be a *n*-tuple. We shall write $\alpha \leq 0$ if $\alpha_i \leq 0$ for all *i*.

Proposition 3.6. Let \mathscr{F} be an exact n-cube of locally free sheaves and, for all $\alpha \leq 0$, let h_{α} be a hermitian metric on \mathscr{F}_{α} . Then there is a unique way to choose metrics h_{α} for all $\alpha \leq 0$, such that $\overline{\mathscr{F}} = \{(\mathscr{F}_{\alpha}, h_{\alpha})\}$ is an emi-n-cube.

Proof. The uniqueness is clear. For the existence, we have to see that, in each \mathscr{F}_{α} , with $\alpha \not\leq 0$, all the possible induced metrics agree. This is guaranteed by the following result.

Lemma 3.7. Let $\{E_{i,j}\}_{i,j} = -1, 0, 1$ be an exact 2-cube of complex vector spaces. Let h be a hermitian metric on $E_{0,0}$ and let $h_{1,0}$ and $h_{0,1}$ be the hermitian metrics in $E_{1,0}$ and $E_{0,1}$ induced by h. Then the metrics induced by $h_{1,0}$ and $h_{0,1}$ in $E_{1,1}$ agree.

Proof. Let us identify $E_{-1,0}$ and $E_{0,-1}$ with their images in $E_{0,0}$. Then the metric $h_{1,0}$ in $E_{1,0}$ is induced by the isomorphism $E_{-1,0}^{\perp} \cong E_{1,0}$. Therefore we can identify $E_{1,0}$ with $E_{-1,0}^{\perp}$ and the morphism $E_{0,0} \longrightarrow E_{1,0}$ with the orthogonal projection. But the image of $E_{0,-1}$ by this orthogonal projection is $(E_{-1,0} + E_{0,-1}) \cap E_{-1,0}^{\perp}$. Therefore the metric in $E_{1,1}$ induced by $h_{1,0}$ is induced by the isomorphism $(E_{-1,0} + E_{0,-1})^{\perp} \cong E_{1,1}$. By symmetry, the same is true for the metric induced by $h_{0,1}$.

(3.5) Let $\mathbb{Z}C_{\text{emi}}(X)$ be the subcomplex of $\mathbb{Z}C(X)$ generated by the emi*n*-cubes, and let D_{emi} be the subcomplex of $\mathbb{Z}C_{\text{emi}}(X)$ generated by the degenerate emi-*n*-cubes. We shall write

$$\widetilde{\mathbb{Z}}C_{\text{emi}}(X) = \mathbb{Z}C_{\text{emi}}(X)/D_{\text{emi}} \subset \widetilde{\mathbb{Z}}C(X)$$

To translate results about emi-*n*-cubes to all exact metrized *n*-cubes we need to construct a morphism of complexes

$$\widetilde{\mathbb{Z}}C(X) \longrightarrow \widetilde{\mathbb{Z}}C_{\text{emi}}(X)$$
.

If $\alpha \in \{-1, 0, 1\}^n$ with $\alpha_i > -1$, we shall write $\alpha - i = (\alpha_1, \dots, \alpha_i - 1, \dots, \alpha_n)$. Let $\overline{\mathscr{F}} = \{(F_\alpha, h_\alpha)\} \in C_n(X)$. For $i = 1, \dots, n$ let $\lambda_i^1 \overline{\mathscr{F}}$ be defined by

$$\lambda_i^1 \overline{\mathscr{F}}_{lpha} = egin{cases} (\mathscr{F}_{lpha}, h_{lpha}), & ext{if } lpha_i = -1, 0, \ (\mathscr{F}_{lpha}, h_{lpha}'), & ext{if } lpha_i = 1 \ , \end{cases}$$

where h'_{α} is the metric induced by $h_{\alpha-i}$. Thus $\lambda_i^1 \overline{\mathscr{F}}$ has the same locally free sheaves as $\overline{\mathscr{F}}$, but we have replaced the metrics of the locally free sheaves of the face $\partial_i^1 \overline{\mathscr{F}}$, by the metrics induced by $\partial_i^0 \overline{\mathscr{F}}$.

Let $\lambda_i^2 \overline{\mathcal{F}}$ be the exact *n*-cube determined by

$$\partial_i^{-1} \lambda_i^2 \overline{\mathscr{F}} = \partial_i^1 \overline{\mathscr{F}},$$

 $\partial_i^0 \lambda_i^2 \overline{\mathscr{F}} = \partial_i^1 \lambda_i^1 \overline{\mathscr{F}},$
 $\partial_i^1 \lambda_i^2 \overline{\mathscr{F}} = 0$.

This *n*-cube measures in some sense the difference between $\overline{\mathcal{F}}$ and $\lambda_i^1 \overline{\mathcal{F}}$. Let us write $\lambda_i \overline{\mathcal{F}} = \lambda_i^1 \overline{\mathcal{F}} + \lambda_i^2 \overline{\mathcal{F}}$, and let us denote by λ the map

 $\lambda_{ij} = \lambda_{ij} + \lambda$

$$\lambda: \mathbb{Z}C_n(X) \longrightarrow \mathbb{Z}C_n(X)$$

$$\overline{\mathscr{F}} \longmapsto \begin{cases} \lambda_n \dots \lambda_1 \overline{\mathscr{F}}, & \text{if } n \ge 1, \\ \overline{\mathscr{F}}, & \text{if } n = 0 \end{cases}$$

Then one can check the following properties:

- (1) λ is a morphism of complexes.
- (2) Im $\lambda \subset \mathbb{Z}C_{\text{emi}}(X)$.
- (3) $\lambda(D) \subset D_{\text{emi}}$.

Therefore this map induces a morphism of complexes

$$\lambda: \widetilde{\mathbb{Z}}C(X) \longrightarrow \widetilde{\mathbb{Z}}C_{\mathrm{emi}}(X)$$

In fact λ is a homotopy equivalence. The inverse being the inclusion $\widetilde{\mathbb{Z}}C_{\text{emi}}(X) \longrightarrow \widetilde{\mathbb{Z}}C(X)$.

(3.6) Let $\overline{\mathscr{F}}$ be an emi-*n*-cube of locally free sheaves. We shall associate to it a locally free sheaf $\operatorname{tr}_n(\overline{\mathscr{F}})$ on $X \times (\mathbb{P}^1)^n$ which, roughly speaking, is a homotopy between the vertexes of $\overline{\mathscr{F}}$.

Let $((x_1:y_1),\ldots,(x_n:y_n))$ be homogeneous coordinates of $(\mathbb{P}^1)^n$. Let \mathscr{I}_{x_i} (resp. \mathscr{I}_{y_i}) be the sheaf of ideals in $X \times (\mathbb{P}^1)^n$ defined by the subvariety $x_i = 0$, (resp. $y_i = 0$). Let $p_0: X \times (\mathbb{P}^1)^n \longrightarrow X$ and $p_i: X \times (\mathbb{P}^1)^n \longrightarrow \mathbb{P}^1$, $i = 1, \ldots, n$, be the projections. Then the maps

$$\begin{split} \mathscr{I}_{x_i} & \xrightarrow{x_i^{-1}} p_i^* \mathcal{O}_{\mathbb{P}^1}(-1) \\ \mathscr{I}_{y_i} & \xrightarrow{y_i^{-1}} p_i^* \mathcal{O}_{\mathbb{P}^1}(-1) \end{split}$$

are isomorphisms. The sheaf $\mathcal{O}_{\mathbb{P}^1}(-1)$ has a metric induced by the standard metric on \mathbb{C}^2 . We put in \mathscr{I}_{x_i} and \mathscr{I}_{y_i} the metrics induced by the above isomorphisms. By 2.9, these metrics are smooth at infinity.

For each pair of integers $i \in \{1, ..., n\}$ and $j \in \{-1, 0\}$, we write

$$\mathscr{I}_{i,j} = \begin{cases} \mathscr{I}_{y_i}, & \text{if } j = -1, \\ \mathscr{I}_{x_i}, & \text{if } j = 0 \end{cases}$$

For each $\alpha \in \langle -1, 0, 1 \rangle^n$, with $\alpha \leq 0$, and for each $k \in \{1, \ldots, n\}$, with $\alpha_k = -1$, we write

$$\begin{split} \mathscr{J}_{\alpha} &= \prod_{i=1}^{n} \mathscr{I}_{i,\alpha_{i}}^{-1} \subset \mathscr{K}_{X \times (\mathbb{P}^{1})^{n}}, \\ \mathscr{J}_{\alpha,k} &= \prod_{i \neq k} \mathscr{I}_{i,\alpha_{i}}^{-1} \subset \mathscr{K}_{X \times (\mathbb{P}^{1})^{n}} \end{split}$$

where $\mathscr{K}_{X \times (\mathbb{P}^1)^n}$ is the sheaf of rational functions on $X \times (\mathbb{P}^1)^n$.

Given an *n*-tuple $\alpha \leq 0$ and an integer $k \in \{1, ..., n\}$, with $\alpha_k = -1$, we write $\alpha + k = (\alpha_1, ..., \alpha_k + 1, ..., \alpha_n)$. We have the inclusions

$$\begin{aligned} \mathcal{J}_{\alpha,k} \subset \mathcal{J}_{\alpha}, \\ \mathcal{J}_{\alpha,k} \subset \mathcal{J}_{\alpha+k} \end{aligned}$$

Let us denote by $\varphi_{\alpha,k} : \overline{\mathscr{F}}_{\alpha} \longrightarrow \overline{\mathscr{F}}_{\alpha+k}$ the morphism $\overline{\mathscr{F}}(\alpha \longrightarrow \alpha + k)$. Let ψ be the morphism

$$\psi:\bigoplus_{\alpha\leq 0} \bigoplus_{k\mid \alpha_{k}=-1} p_{0}^{*\overline{\mathscr{F}}_{\alpha}} \otimes \mathscr{J}_{\alpha,k} \longrightarrow \bigoplus_{\alpha\leq 0} p_{0}^{*\overline{\mathscr{F}}_{\alpha}} \otimes \mathscr{J}_{\alpha+k} \ ,$$

which sends $s \otimes g \in p_0^* \overline{\mathscr{F}}_{\alpha} \otimes \mathscr{J}_{\alpha,k}$ to

$$egin{aligned} \psi(s\otimes g) &= s\otimes g + arphi_{lpha,k}(s)\otimes g \ &\in p_0^*\overline{\mathscr{F}}_{lpha}\otimes \mathscr{J}_{lpha} \oplus p_0^*\overline{\mathscr{F}}_{lpha+k}\otimes \mathscr{J}_{lpha+k} \end{aligned}$$

The locally free sheaf $\bigoplus_{\alpha \leq 0} p_0^* \overline{\mathscr{F}}_{\alpha} \otimes \mathscr{J}_{\alpha}$ has a metric induced by the metrics of \mathscr{I}_{x_i} , \mathscr{I}_{y_i} and $\overline{\mathscr{F}}_{\alpha}$. This metric is smooth at infinity.

Definition 3.8. The *n*-transgression of $\overline{\mathcal{F}}$ is the hermitian locally free sheaf

$$\operatorname{tr}_n(\overline{\mathscr{F}}) = \operatorname{Coker}(\psi)$$
,

with the metric induced by the metric of $\bigoplus_{\alpha \leq 0} p_0^* \mathscr{F}_{\alpha} \otimes \mathscr{J}_{\alpha}$. By Proposition 2.8, this metric is smooth at infinity.

The following result follows directly from the definition.

Proposition 3.9. Let $\overline{\mathcal{F}}$ be an emi-*n*-cube. Then there are isometries

$$\begin{aligned} \operatorname{tr}_{n}(\overline{\mathscr{F}})\big|_{\{x_{i}=0\}} &\cong \operatorname{tr}_{n-1}\left(\partial_{i}^{0}\overline{\mathscr{F}}\right) ,\\ \operatorname{tr}_{n}(\overline{\mathscr{F}})\big|_{\{y_{i}=0\}} &\cong \operatorname{tr}_{n-1}\left(\partial_{i}^{-1}\overline{\mathscr{F}}\right) \stackrel{\perp}{\oplus} \operatorname{tr}_{n-1}\left(\partial_{i}^{1}\overline{\mathscr{F}}\right) \end{aligned}$$

(3.7) Let us give an inductive construction of the transgressions. If n = 1, an emi-1-cube, $\overline{\mathscr{F}}$ is a short exact sequence

$$\overline{\mathscr{F}}_{-1} \xrightarrow{f} \overline{\mathscr{F}}_{0} \longrightarrow \overline{\mathscr{F}}_{1} \quad ,$$

where the metric of $\overline{\mathscr{F}}_1$ is induced by the metric of $\overline{\mathscr{F}}_0$. Then $\operatorname{tr}_1(\overline{\mathscr{F}})$ is the cokernel of the map

$$\begin{array}{ccc} \mathscr{F}_{-1} \longrightarrow \mathscr{F}_{-1} \otimes \mathscr{I}_{y_{1}}^{-1} \oplus \mathscr{F}_{0} \otimes \mathscr{I}_{x_{1}^{-1}} \\ s \longmapsto & s \otimes 1 & \oplus f(s) \otimes 1 \end{array} .$$

Observe that this is a minor modification of the locally free sheaf used by Bismut, Gillet and Soulé [B-G-S], [G-S 1] to construct Bott-Chern forms. In the definition given here, we avoid the use of partitions of unity, obtaining a natural construction. The price is to restrict ourselves to emi-*n*-cubes.

If $\overline{\mathscr{F}}$ is an emi-*n*-cube, let $\operatorname{tr}_1(\overline{\mathscr{F}})$ be the emi-*n* - 1-cube over $X \times \mathbb{P}^1$ defined by:

$$\operatorname{tr}_1(\overline{\mathscr{F}})_{\alpha} = \operatorname{tr}_1(\partial_{n^{\alpha}}^{\alpha}\overline{\mathscr{F}})$$

Then we write

$$\operatorname{tr}_k(\overline{\mathscr{F}}) = \operatorname{tr}_1(\operatorname{tr}_{k-1}(\overline{\mathscr{F}}))$$

The hermitian locally free sheaf $tr_n(\overline{\mathscr{F}})$ defined in this way coincides with the earlier definition. Thus the transgressions are simply an iteration of the construction of Bismut, Gillet and Soulé.

(3.8) For any homology complex A_* , we shall denote by A^* the cohomology complex defined by $A^k = A_{-k}$. Let us use the transgressions previously defined, to associate to every emi-*n*-cube a family of differential forms.

Definition 3.10. Let

$$ch: \mathbb{Z}C^*_{emi}(X) \longrightarrow \bigoplus_p \widetilde{\mathfrak{H}}^*(X,p)[2p]$$

be the map given by

$$\operatorname{ch}(\overline{\mathscr{F}}) = \widetilde{\operatorname{ch}}_0(\operatorname{tr}_n(\overline{\mathscr{F}}))_{\mathscr{H}}$$
,

where $\widetilde{ch}_0(\cdot)_{\mathscr{H}}$ is as in (2.4).

Proposition 3.11. The map ch is a morphism of complexes and factorizes through a unique morphism

$$ch: \widetilde{\mathbb{Z}}C^*_{\mathrm{emi}}(X) \longrightarrow \bigoplus_p \widetilde{\mathfrak{H}}^*(X,p)[2p]$$

Proof. To see that it is a morphism of complexes, observe that, since the forms $\widetilde{ch}_0(\cdot)_{\mathscr{H}}$ are closed,

$$dch(tr_n(\overline{\mathscr{F}})) = \sum_{i=1}^n \sum_{j=0}^{1} (-1)^{i+j} d_i^j \widetilde{ch}_0(tr_n(\overline{\mathscr{F}}))_{\mathscr{H}}$$
$$= \sum_{i=1}^n (-1)^i \widetilde{ch}_0\left(tr_n\left(\overline{\mathscr{F}}\right)\right)_{\mathscr{H}} \Big|_{\{x_i=0\}} + \sum_{i=1}^n (-1)^{i+1} \widetilde{ch}_0\left(tr_n\left(\overline{\mathscr{F}}\right)\right)_{\mathscr{H}} \Big|_{\{y_i=0\}}$$
$$= \sum_{i=1}^n (-1)^i \widetilde{ch}_0\left(tr_n\left(\overline{\mathscr{F}}\right)\Big|_{\{x_i=0\}}\right)_{\mathscr{H}} + \sum_{i=1}^n (-1)^{i+1} \widetilde{ch}_0\left(tr_n\left(\overline{\mathscr{F}}\right)\Big|_{\{y_i=0\}}\right)_{\mathscr{H}}$$

Therefore, by Proposition 3.9,

$$d\mathrm{ch}(\mathrm{tr}_n(\overline{\mathscr{F}})) = \sum_{i=1}^n \sum_{j=-1}^1 (-1)^{i+j} \widetilde{\mathrm{ch}}_0(\mathrm{tr}_{n-1}(\partial_i^j \overline{\mathscr{F}}))_{\mathscr{H}}$$

 $= \mathrm{ch}(d\overline{\mathscr{F}}) \; .$

To see the existence of the factorization, we have to show that, for a degenerate emi-*n*-cube $\overline{\mathscr{F}}$, we have $\operatorname{ch}(\overline{\mathscr{F}}) = 0$ in $\bigoplus \widetilde{\mathfrak{H}}^*(X,p)$. By symmetry we may assume that $\overline{\mathscr{F}} = s_n^j \mathscr{G}$, with $j \in \{-1, 1\}$ and $\overline{\mathscr{G}}$ an emi*n*-1-cube.

If j = 1, then $\operatorname{tr}_{n-1}(\overline{\mathcal{F}})$ is the exact sequence

$$0 \longrightarrow (\mathrm{Id} \times s^n)^* \mathrm{tr}_{n-1}(\overline{\mathscr{G}}) \xrightarrow{\mathrm{Id}} (\mathrm{Id} \times s^n)^* \mathrm{tr}_{n-1}(\overline{\mathscr{G}}) \longrightarrow 0 \longrightarrow 0 \,.$$

Therefore $\operatorname{tr}_n(\overline{\mathcal{F}})$ is the cokernel of the map

$$\begin{array}{c} (\mathrm{Id} \times s^n)^* \mathrm{tr}_{n-1}(\overline{\mathscr{G}}) \longrightarrow (\mathrm{Id} \times s^n)^* \mathrm{tr}_{n-1}(\overline{\mathscr{G}}) \otimes \mathscr{I}_{y_n}^{-1} \oplus (\mathrm{Id} \times s^n)^* \mathrm{tr}_{n-1}(\overline{\mathscr{G}}) \otimes \mathscr{I}_{x_n}^{-1} \\ x \longmapsto x \otimes 1 \quad \oplus \quad x \otimes 1 \quad . \end{array}$$

But $\mathscr{I}_{y_n}^{-1}$ and $\mathscr{I}_{x_n}^{-1}$ are both isometric with $p_n^*\mathcal{O}(1)$. Hence this cokernel is isometric with $(\mathrm{Id} \times s^n)^* \mathrm{tr}_{n-1}(\overline{\mathscr{G}}) \otimes p_n^*\mathcal{O}(2)$, where $\mathcal{O}(2)$ is provided with the standard metric. Thus

$$\widetilde{\mathrm{ch}}_{0}(\mathrm{tr}_{n}(\overline{\mathscr{F}}))_{\mathscr{H}} = (\mathrm{Id} \times s^{n})^{*} \widetilde{\mathrm{ch}}_{0}(\mathrm{tr}_{n-1}(\overline{\mathscr{G}}))_{\mathscr{H}} + 2\omega_{n} \wedge (\mathrm{Id} \times s^{n})^{*} \widetilde{\mathrm{ch}}_{0}(\mathrm{tr}_{n-1}(\overline{\mathscr{G}}))_{\mathscr{H}}.$$

which is zero in $\bigoplus \widetilde{\mathfrak{H}}^*(X, p)$.

The case j = -1 is analogous.

Definition 3.12. We shall denote also by ch the composition

$$\widetilde{\mathbb{Z}}C^*(X) \xrightarrow{\lambda} \widetilde{\mathbb{Z}}C^*_{\operatorname{emi}}(X) \xrightarrow{\operatorname{ch}} \bigoplus_p \widetilde{\mathfrak{H}}^*(X,p)[2p]$$

Definition 3.13. Let $\overline{\mathcal{F}}$ be a metrized exact *n*-cube. The form $\operatorname{ch}(\lambda(\overline{\mathcal{F}}))$ will be called the Bott-Chern form of $\overline{\mathcal{F}}$ and will be denoted by $\operatorname{ch}_{n}(\overline{\mathcal{F}})_{\mathscr{H}}$.

4 Higher characteristic classes

The Chern character from *K*-theory to a suitable cohomology theory, such as absolute Hodge cohomology, is additive for exact sequences. Nevertheless, given a cochain complex which computes absolute Hodge cohomology, we cannot make a consistent choice of representatives of the Chern character that behaves additively. Following the ideas of Schechtman ([Sch]), the lack of additivity at the level of complexes, of the Chern character for K_n , gives us the Chern character for K_{n+1} .

In the previous section we have associated, to each metrized exact *n*-cube, a family of differential forms. The differential form associated to an *n*-cube measures the lack of additivity of the differential forms associated to its faces. In this section we shall see that this construction allows us to define higher Chern character classes from *K*-theory to absolute Hodge cohomology.

(4.1) Let us begin by reviewing the Waldhausen *K*-theory of a small exact category. We shall follow [Sch] (See also [Wal] or [Lo 1]).

For $n \in \mathbb{N}$, let Cat(n) denote the category associated with the ordered set $\{1, \ldots, n\}$. Let M_n be the category of morphisms of Cat(n). That is

Ob
$$M_n = \{(i, j) \in \mathbb{N} \times \mathbb{N} \mid 0 \le i \le j \le n\}$$

and Hom((i, j), (k, l)) contains a unique element if $i \le k$ and $j \le l$ and is empty otherwise. The categories M_n form a cosimplicial category M.

For any category \mathfrak{C} , let us denote by $M_n\mathfrak{C}$ the category of functors from M_n to \mathfrak{C} .

Definition 4.1. Let \mathfrak{E} be a small exact category and 0 a fixed zero object of \mathfrak{E} . Let $\underline{S}_n \mathfrak{E}$ be the full subcategory of $M_n \mathfrak{E}$, whose objects are the functors $M_n \longrightarrow \mathfrak{E}$, such that,

(1) for all $i, E_{i,i} = 0$; (2) for all $i \le j \le k$,

$$E_{i,j} \longrightarrow E_{i,k} \longrightarrow E_{j,k}$$

is a short exact sequence.

Let us write $S_n \mathfrak{E} = \operatorname{Ob} \underline{S}_n \mathfrak{E}$. We shall denote by $\underline{S}\mathfrak{E}$ or $\underline{S}^1\mathfrak{E}$ the simplicial exact category $\amalg \underline{S}_n \mathfrak{E}$, and by $S\mathfrak{E}$ or $S^1\mathfrak{E}$ the simplicial set $\operatorname{Ob} \underline{S}\mathfrak{E}$.

In other words we have:

$$S_{0} \mathfrak{E} = \{0\},$$

$$S_{1} \mathfrak{E} = \text{Ob } \mathfrak{E},$$

$$S_{2} \mathfrak{E} = \{\text{exact sequences of } \mathfrak{E}\},$$

$$S_{n} \mathfrak{E} = \begin{cases} \text{sequences of monomorphisms} \\ E_{0,1} \longrightarrow E_{0,2} \longrightarrow \cdots \longrightarrow E_{0,n} \\ \text{with a choice of quotients} \\ E_{i,j} \cong E_{0,j}/E_{0,i} \end{cases} \end{cases}$$

In particular $S\mathfrak{E}$ is a pointed simplicial set. In the sequel we shall sometimes use the word space to denote simplicial sets.

For a space C, we shall denote by |C| its geometric realization.

Proposition 4.2. (Cf. [Lo 2].) There is a homotopy equivalence

$$S\mathfrak{E} \cong BQ\mathfrak{E}$$
,

where Q denotes Quillen's Q-construction and B means classifying space. Therefore, for all $i \ge 0$,

$$K_i(\mathfrak{E}) = \pi_{i+1}(|S_{\cdot}\mathfrak{E}|, 0)$$
.

(4.2) Let us recall the notion of spectrum from [Th]. For any pointed space C, let us write ΣC for the suspension of C, and ΩC for the loop space of C. We shall use the same notation for topological spaces.

Definition 4.3. A prespectrum X is a sequence of pointed spaces X_n for all nonnegative integers n, together with structure maps $\Sigma X_n \longrightarrow X_{n+1}$. These maps can also be described by their adjoint $X_n \longrightarrow \Omega X_{n+1}$. A fibrant spectrum is a prespectrum such that all X_n are fibrant spaces and the structure maps $X_n \longrightarrow \Omega X_{n+1}$ are weak equivalences.

The space $S\mathfrak{E}$ is a piece of a prespectrum. To construct the other spaces that form the prespectrum, we write inductively

$$\underline{S}^{m} \mathfrak{E} = \underline{SS}^{m-1} \mathfrak{E},$$
$$S^{m} \mathfrak{E} = S \underline{S}^{m-1} \mathfrak{E}.$$

Then \underline{S}^m is an exact m-simplicial category and \underline{S}^m is a m-simplicial set. For a poly-simplicial set C let diag(C) denote its diagonal space. We shall denote by |C| = |diag(C)| its geometric realization.

Proposition 4.4. (Schechtman [Sch 1.2].) There are natural maps

$$\varphi_m: \Sigma S^m \mathfrak{E} \longrightarrow S^{m+1} \mathfrak{E}$$
,

inducing homotopy equivalences

$$|S^m\mathfrak{E}|\cong \Omega |S^{m+1}\mathfrak{E}|$$

As a consequence of this proposition, if we write $S^0 \mathfrak{E} = \Omega S \mathfrak{E}$, then the sequence of spaces diag($S^m \mathfrak{E}$) is a prespectrum. Moreover, if we replace the above spaces by weakly equivalent fibrant spaces we shall obtain a fibrant spectrum. For instance, let us denote by Sing the singular functor (see [B-K]). Then, if we write

$$\mathbf{K}_m(\mathfrak{E}) = \operatorname{Sing}(|S^m\mathfrak{E}|) \;\;,$$

the spaces \mathbf{K}_m form a fibrant spectrum. By Proposition 4.2, the homotopy of this fibrant spectrum is the *K*-theory of \mathfrak{E} .

(4.3) For example, let X be a smooth complex variety, let $m \ge 1$ be an integer and let us write $S^m(X) = S^m(\overline{\mathfrak{E}}(X))$. Then, for $i \ge 0$, the *i*-th K-groups of X is

$$K_i(X) = \pi_{i+m}(|S^m(X)|, 0)$$
.

By Proposition 4.4 this definition does not depend on the choice of m.

(4.4) Let us associate, to each element of $S_n \mathfrak{E}$, an exact n - 1-cube. We shall do so inductively. For n = 1, we write

$$\operatorname{Cub}(\{E_{i,j}\}_{0 \le i \le j \le 1}) = E_{0,1}$$
.

Assume that we have defined Cub *E* for all $E \in S_m \mathfrak{E}$, with m < n. Let $E \in S_n \mathfrak{E}$. Then Cub *E* is the n - 1-cube with

$$\partial_1^{-1} \operatorname{Cub} E = s_{n-2}^1 \dots s_1^1(E_{0,1}),$$

$$\partial_1^0 \operatorname{Cub} E = \operatorname{Cub} (\partial_1 E),$$

$$\partial_1^{-1} \operatorname{Cub} E = \operatorname{Cub} (\partial_0 E) .$$

For instance, if n = 2, then Cub $(\{E_{i,j}\}_{0 \le i \le j \le 2})$ is the short exact sequence

$$E_{0,1} \longrightarrow E_{0,2} \longrightarrow E_{1,2}$$

On the other hand, if n = 3, then $\text{Cub}(\{E_{i,j}\}_{0 \le i \le j \le 3})$ is the exact square

$E_{0,1}$	\longrightarrow	$E_{0,2}$	\longrightarrow	$E_{1,2}$
_↓				_
$E_{0,1}$	\longrightarrow	$E_{0,3}$	\longrightarrow	$E_{1,3}$
Ļ		_↓		_↓
0	\longrightarrow	$E_{2,3}$	\longrightarrow	$E_{2,3}$

All the faces of the n - 1-cube Cub E can be computed explicitly.

Proposition 4.5. Let $E \in S_n \mathfrak{E}$. Then, for i = 1, ..., n - 1, the faces of the n - 1-cube Cub E are

$$\partial_i^{-1} \operatorname{Cub} E = s_{n-2}^1 \dots s_i^1 \operatorname{Cub} \partial_{i+1} \dots \partial_n E,$$

$$\partial_i^0 \operatorname{Cub} E = \operatorname{Cub} \partial_i E,$$

$$\partial_i^1 \operatorname{Cub} E = s_{i-1}^{-1} \dots s_1^{-1} \operatorname{Cub} \partial_0 \dots \partial_{i-1} E.$$

By proposition 4.5 and using induction we have,

Corollary 4.6. The n - 1-cube Cub E is exact.

Therefore we have a map $\operatorname{Cub} : S_n \mathfrak{E} \longrightarrow C_{n-1} \mathfrak{E}$.

(4.5) Let $\mathbb{Z}S\mathfrak{E}$ be the homological complex associated with the simplicial set $S\mathfrak{E}$. That is, $\mathbb{Z}S_n\mathfrak{E}$ is the free abelian group generated by $S_n\mathfrak{E}$, and the differential $d : \mathbb{Z}S_n\mathfrak{E} \longrightarrow \mathbb{Z}S_{n-1}\mathfrak{E}$ is given by

$$d = \sum_{i=0}^n (-1)^i \partial_i$$
 .

The map Cub can be extended by linearity to a map

$$\operatorname{Cub}: \mathbb{Z}S\mathfrak{E}[1] \longrightarrow \mathbb{Z}C\mathfrak{E}$$
.

Note that this map is not a morphism of complexes. However, the map Cub induces a map also denoted by Cub : $\mathbb{Z}S\mathfrak{E}[1] \longrightarrow \widetilde{\mathbb{Z}}C\mathfrak{E}$. And, since by Proposition 4.5,

dCub E = Cub dE + degenerate elements ,

we have:

Corollary 4.8. The map $\operatorname{Cub} : \mathbb{Z}S\mathfrak{E}[1] \longrightarrow \widetilde{\mathbb{Z}}C\mathfrak{E}$ is a morphism of complexes.

(4.6) We can obtain analogous maps for all the spaces $S^m \mathfrak{E}$. In particular, we have maps

 $\operatorname{Cub}: S_{n_1} \dots S_{n_m} \mathfrak{E} \longrightarrow C_{n_1-1} \dots C_{n_m-1} \mathfrak{E} \longrightarrow C_{n_1+\dots+n_m-m} \mathfrak{E}$.

Let us denote by $\mathbb{Z}S^m \mathfrak{E}$ the chain complex that, in degree *n*, is the free abelian group generated by

$$\prod_{n_1+\cdots+n_m=n}S_{n_1}\ldots S_{n_m}\mathfrak{E}$$

The differential of this complex is the alternate sum of all the face maps. Note that this complex is homotopically equivalent to \mathbb{Z} diag ($S^m \mathfrak{E}$). The induced map

J. I. Burgos, S. Wang

$$\operatorname{Cub}: \mathbb{Z}S^m\mathfrak{E}[m] \longrightarrow \mathbb{Z}C\mathfrak{E}$$

is also a morphism of complexes.

(4.7) Let $m \ge 1$ be an integer. We shall denote by $\mathbb{Z}S_m^*(X)$ the cohomological complex associated to the homological complex $\mathbb{Z}S_m^*(X)$.

Definition 4.9. The Chern character map is the composition

$$\mathbb{Z}S^*_m(X)[-m] \xrightarrow{\operatorname{Cub}} \widetilde{\mathbb{Z}}C^*(X) \xrightarrow{\lambda} \widetilde{\mathbb{Z}}C^*_{\operatorname{emi}}(X) \xrightarrow{\operatorname{ch}} \bigoplus_p \widetilde{\mathfrak{F}}^*(X,p)[2p] .$$

This map will also be denoted by ch. The Chern character classes are obtained by composing with the Hurewick map:

$$K_i(X) = \pi_{i+m}(S^m(X)) \longrightarrow H_{i+m}(\mathbb{Z}S^m(X)) \longrightarrow \bigoplus_p H^{2p-i}_{\mathscr{H}}(X,p) \quad .$$

Proposition 4.10. The above definition does not depend on the choice of m.

Proof. For a pointed simplicial set S, with base point p, we shall write

$$\mathbb{Z}'S_*=\mathbb{Z}S_*/\mathbb{Z}p_*$$
 .

The natural map $\Sigma S^m(X) \longrightarrow S^{m+1}(X)$ induces a morphism

$$\mathbb{Z}'S^m_*(X)[m] \longrightarrow \mathbb{Z}S^{m+1}_*(X)[m+1]$$

By the proof of Proposition 4.4 in [Sch 1.2] this map is induced by the natural bijection $S^m(X) \longrightarrow S_1 S^m(X)$. Therefore the diagram



is commutative. By the commutativity of this diagram and proposition 4.4 the Chern character classes are independent of the choice of m

5 Beilinson's regulator

The aim of this section is to prove that the higher Chern character classes defined in Sect. 4 agree with Beilinson's regulator map.

(5.1) Let us begin by extending the definition of the map ch to the case of simplicial smooth complex varieties. To this end, we first recall the con-

struction of the absolute Hodge cohomology of X = X. a smooth simplicial complex variety. For each p, the complexes $\tilde{\mathfrak{H}}^*(X_n, p)$ form a cosimplicial complex as n varies. Let $\mathscr{N}\mathfrak{H}^*(X, p)$ be the associated double complex and let us denote the simple complex by

$$\widetilde{\mathfrak{H}}^*(X,p) = s(\mathscr{N}\widetilde{\mathfrak{H}}^*(X,p))$$
.

Then

$$H^*_{\mathscr{H}}(X, \mathbb{R}(p)) = H^*(\mathfrak{H}^*(X, p))$$

For the definition of the *K*-theory of simplicial schemes we shall follow [Sch]. We shall say that a smooth simplicial scheme X = X has finite dimension if there is an integer *m* such that

$$X = \operatorname{Sk}_m(X)$$
,

where $Sk_m(X)$ is the *m*-th skeleton of *X*, that is, the simplicial scheme generated by X_0, \ldots, X_m .

Let X = X be a simplicial scheme of finite dimension. The family of prespectrums $\{S(X_n)\}_n$ form a cosimplicial prespectrum S(X). Let $\mathbf{K}(X)$ be a fibrant cosimplicial fibrant spectrum weakly equivalent to S(X). Then the K groups of X are defined as

$$K_i(X) = \pi_{i+1}(\text{Tot } \mathbf{K}(X))$$

Since X has finite dimension, there is a convergent spectral sequence

$$E_1^{p,q} = K_{-q}(X_p) \Longrightarrow K_{-p-q}(X_{\cdot})$$
.

Observe that for a given simplicial scheme X., of finite dimension, it is not necessary to work with the whole spectrum. Let m be such that $X = \text{Sk}_m X$. Let us choose a positive integer q, and let $\mathbf{K}_q(X)$ be a fibrant cosimplicial fibrant space, weakly equivalent to $S^q(X)$. If q > m or q > -i, then

$$K_i(X_{\cdot}) = \pi_{i+q}(\text{Tot } \mathbf{K}_q(X_{\cdot}))$$
.

For an arbitrary simplicial scheme we write

$$\widehat{K}_*(X_{\cdot}) = \lim_{\stackrel{\longleftarrow}{m}} K_*(\operatorname{Sk}_m(X_{\cdot}))$$
.

Let X be a smooth simplicial complex variety of finite dimension. Since the map ch defined in Sect. 4 gives us a morphism of complexes

$$ch: s\mathcal{N}\mathbb{Z}S_q^*(X)[q] \longrightarrow \bigoplus_p \widetilde{\mathfrak{H}}^*(X,p)[2p] ,$$

we can extend the definition of the Chern character to the simplicial case, obtaining maps:

$$\operatorname{ch}: K_i(X) \longrightarrow \bigoplus_p H^{2p-i}_{\mathscr{H}}(X, \mathbb{R}(p))$$

If X does not have finite dimension, taking limits, we have also characteristic classes

$$ch: \widehat{K}_i(X) \longrightarrow \bigoplus_p H^{2p-i}_{\mathscr{H}}(X, \mathbb{R}(p))$$

Remark. 5.1. All the constructions needed to define the map ch can be extended to the case of a smooth simplicial scheme X. over \mathbb{C} , such that each X_n is a (not necessarily finite) disjoint union of smooth complex varieties. For instance, by a compactification of X_n we shall mean a disjoint union of compactifications of each component of X_n .

(5.2) Beilinson [Be] has defined characteristic classes from K-theory to absolute Hodge cohomology. These classes are a particular case of the characteristic classes defined by Gillet [Gi] to any suitable cohomology theory. In particular, Beilinson's regulator is the Chern character in this theory. Let us denote by ρ the Beilinson's regulator map.

Then ch and ρ are natural transformations between contravariant functors. Both agree with the classical Chern character on the K_0 groups of smooth complex varieties. The aim of this section is to prove the following theorem.

Theorem 5.2. Let X be a smooth complex variety. Let $\sigma \in K_i(X)$. Then $ch(\sigma) = \rho(\sigma)$.

Proof. Let $\mathfrak{U} = \{U_{\alpha}\}$ be an open covering of X. We shall denote by $\mathfrak{E}(X, \mathfrak{U})$ the full subcategory of $\mathfrak{E}(X)$ composed by the locally free sheaves on X whose restrictions to all U_{α} are free. We shall denote by $\overline{\mathfrak{E}}(X, \mathfrak{U})$ the category of hermitian vector bundles on X whose restrictions to all U_{α} are free. Let us write

$$K_i(X, \mathfrak{U}) = \pi_{i+1}(S\mathfrak{E}(X, \mathfrak{U})) = \pi_{i+1}(S\overline{\mathfrak{E}}(X, \mathfrak{U}))$$
.

Then

$$K_i(X) = \varinjlim_{\mathfrak{U}} K_i(X,\mathfrak{U})$$
.

(5.2.1) Following Schechtman ([Sch]) we know that there is a simplicial scheme \mathscr{BP} , which is a classifying space for algebraic *K*-theory. More precisely, Schechtman proves the following result.

Theorem 5.3. (Schechtman) *There is a homotopy equivalence*

$$S\mathfrak{E}(X,\mathfrak{U}) \cong \underline{\mathrm{Hom}}(N\mathfrak{U},\mathscr{B}P)$$

where <u>Hom</u> is the function space and $N\mathfrak{U}$ is the nerve of the covering.

(5.2.2) Let Y = Y be a smooth simplicial scheme of finite dimension. Let us denote by $\underline{\text{Hom}}(Y, \mathscr{B}P)$ the cosimplicial simplicial set $\underline{\text{Hom}}(Y, \mathscr{B}P)_m^n$ = $\text{Hom}(Y_n, \mathscr{B}P_m)$. Then

$$\operatorname{Hom}(Y, \mathscr{B}P) = \operatorname{Tot} \operatorname{Hom}(Y, \mathscr{B}P)$$
.

For any scheme X, let σX be the simplicial scheme with $\sigma X_n = X$ and all the faces and degeneracies equal to the identity. Then the simplicial set Hom $(X, \mathscr{B}P)$ is the function space Hom $(\sigma X, \mathscr{B}P)$. Observe that σX is the nerve of the trivial covering $\{X\}$. Thus, by Theorem 5.3 and the comparison between Tot and Holim (cf. [B-K, XI, 4.4], [Th, 5.25] and [Le, 3.1.2]) we obtain a natural map

$$\operatorname{Tot} \operatorname{\underline{Hom}}(Y, \mathscr{B}P) \longrightarrow \operatorname{Holim} S\mathfrak{E}(Y_n, \{Y_n\}) \quad .$$

Taking homotopy groups we obtain natural maps

$$\pi_i \underline{\operatorname{Hom}}(Y, \mathscr{B}P) \longrightarrow K_{i-1}(Y)$$
.

In particular, if $f: Y \longrightarrow \mathscr{B}P$ is simplical morphism of simplicial schemes, then f defines an element of $\pi_0 \underline{\text{Hom}}(Y, \mathscr{B}P)$. Let us denote by e_f the image of this element in $K_{-1}(Y)$.

If Y does not have finite dimension, since any simplicial morphism $f: Y \longrightarrow \mathscr{B}P$ induces simplicial morphisms $f_m: \operatorname{Sk}_m Y \longrightarrow \mathscr{B}P$, taking a limit, we obtain an element $e_f \in \widehat{K}_{-1}(Y)$.

Remark 5.4. The identity of $\mathscr{B}P$ defines an element, denoted by $e_{\mathscr{B}P} \in \widehat{K}_{-1}(\mathscr{B}P)$. Moreover, for f as above, $e_f = f^*(e_{\mathscr{B}P})$.

(5.2.3) The element e_{gp} is, in some sense, a universal element in *K*-theory. Since $e_{gp} \in \widehat{K}_{-1}$, to exploit the universality of this element, we need to relate elements in \widehat{K}_n with elements in \widehat{K}_{-1} . This can be done using spheres.

Let $\sigma \in K_n(X)$. Then there is an open covering \mathfrak{U} of X, such that $\sigma \in K_n(X, \mathfrak{U}) = \pi_{n+1}(\underline{\operatorname{Hom}}(N\mathfrak{U}, \mathscr{B}P))$. Therefore, there exists an integer $d \ge 0$ such that σ is represented by an element

$$\gamma_{\sigma} \in \underline{\operatorname{Hom}}(\operatorname{Sub}_{d} S^{n+1} \times N\mathfrak{U}, \mathscr{B}P) = \underline{\operatorname{Hom}}(\operatorname{Sub}_{d} S^{n+1}, \underline{\operatorname{Hom}}(N\mathfrak{U}, \mathscr{B}P))$$

where S^{n+1} is the (pointed) simplicial n + 1-dimensional sphere and Sub_d is the *d*-th subdivision. Let us denote by $\sum_{n+1}^{n+1} = \operatorname{Sub}_d S^{n+1}$.

Lemma 5.5. Let Y = Y be a smooth simplicial complex variety. Then there are *natural decompositions*

$$\widehat{K}_{-1}(\sum^{n+1} \times Y) = \widehat{K}_{-1}(Y) \oplus \widehat{K}_n(Y),$$
$$H^{2p+1}_{\mathscr{H}}\left(\sum^{n+1} \times Y, \mathbb{R}(p)\right) = H^{2p+1}_{\mathscr{H}}(Y, \mathbb{R}(p)) \oplus H^{2p-n}_{\mathscr{H}}(Y, \mathbb{R}(p))$$

Moreover, the maps ch and ρ are compatible with these decompositions.

Proof. We may assume that *Y* has finite dimension because the general case is obtained taking the limit. Then

$$K_{-1}\left(\sum_{\alpha}^{n+1} \times Y\right) = \pi_0(\operatorname{Tot}_{\alpha}(\operatorname{Tot}_{\beta}(\mathbf{K}(\sum_{\alpha}^{n+1} \times Y_{\beta}))))$$

The spectral sequence associated with Tot_{α} has E_2 -term:

$$E_2^{p,q} = \begin{cases} K_{-q}(Y), & \text{if } p = 0, n+1, \\ 0, & \text{if } p \neq 0, n+1 \end{cases}$$

Let us denote by * the simplicial point. Since the spectral sequence of $* \times Y$ splits the spectral sequence of $\sum^{n+1} \times Y$, the above spectral sequence degenerates at the E_2 -term, and the exact sequence obtained from this spectral sequence splits in a natural way.

The same argument works for cohomology. Moreover, since ch and ρ are natural transformations, they induce morphisms between the *K*-theoretical and the cohomological spectral sequences, proving the compatibility statement.

Let us denote by $pr: K_{-1}(\sum^{n+1} \times N\mathfrak{U}) \longrightarrow K_n(N\mathfrak{U})$ the projection. The precise meaning of the universality of $e_{\mathscr{B}P}$ is given by the following result.

Lemma 5.6. In the group $K_n(N\mathfrak{U})$, the equality

$$\operatorname{pr}(\gamma_{\sigma}^*(e_{\mathscr{B}P})) = \sigma$$

holds.

Proof. By Remark 5.4,

$$\operatorname{pr}(\gamma_{\sigma}^{*}(e_{\mathscr{B}P})) = \operatorname{pr}(e_{\gamma_{\sigma}})$$
.

On the other hand, by the definition of γ_{σ} , the map

$$\pi_0(\underline{\operatorname{Hom}}(\sum^{n+1} \times N\mathfrak{U}, \mathscr{B}P)) \xrightarrow{\operatorname{pr}} \pi_{n+1}(\underline{\operatorname{Hom}}(N\mathfrak{U}, \mathscr{B}P))$$

sends the class of γ_{σ} to the class of σ . Therefore, since the diagram

$$\begin{array}{ccc} \pi_0(\underbrace{\operatorname{Hom}}_{\operatorname{pr}\downarrow} \times N\mathfrak{U}, \mathscr{B}P)) & \longrightarrow & K_{-1}(\sum^{n+1} \times N\mathfrak{U}) \\ & & & & \\ \pi_{n+1}(\underbrace{\operatorname{Hom}}_{n+1}(\mathcal{N}\mathfrak{U}, \mathscr{B}P)) & \longrightarrow & K_n(N\mathfrak{U}) \end{array} .$$

is commutative we have that

$$\operatorname{pr}\gamma_{\sigma}^{*}(e_{\mathscr{B}P})) = \sigma$$
.

(5.2.4) By Remark 5.1, the map ch is defined for the simplicial scheme \mathcal{BP} . Moreover, by the naturality of ch and ρ and their compatibility with the map pr, we have

$$ch(\sigma) = pr(\gamma^*_{\sigma}(ch(e_{\mathscr{B}P}))), \rho(\sigma) = pr(\gamma^*_{\sigma}(\rho(e_{\mathscr{B}P}))) .$$

Thus, to prove Theorem 5.2, we are led to compare $ch(e_{\mathscr{R}P})$ and $\rho(e_{\mathscr{R}P})$. For this comparison, we need to understand the cohomology of $\mathscr{R}P$. This cohomology has been computed by Schechtman [Sch]. The simplicial scheme $\mathscr{R}P$ is the classifying space of a simplicial group P, where $P_0 = *$ and $P_1 = \prod_n GL(n)$. Thus it is a bisimplicial scheme $\mathscr{R}.P$. The edge homomorphism of the spectral sequence associated to the second index gives us a morphism

$$d_{H}: H^{2p+1}_{\mathscr{H}}(\mathscr{B}P, \mathbb{R}(p)) \longrightarrow \prod_{n \ge 0} H^{2p}_{\mathscr{H}}(\mathrm{BGL}(n), \mathbb{R}(p)) \quad .$$

Let us denote by $A = H^*_{\mathscr{H}}(\operatorname{Spec} \mathbb{C}, \mathbb{R}(*)).$

For each i, n let us denote by

$$c_{i,n} = c_i(E_n) \in H^{2i}_{\mathscr{H}}(\mathrm{BGL}(n), \mathbb{R}(i))$$

the *i*-th Chern class of the tautological vector bundle over BGL(n). Then we have an isomorphism

$$H^*_{\mathscr{H}}(\mathrm{BGL}(n),\mathbb{R}(*))=A[c_{1,n},\ldots,c_{n,n}]$$
.

Let $s_{k,n} \in A[c_{1,n}, \ldots, c_{n,n}]$ be the *k*-th Newton polynomial in the $c_{i,n}$. That is, $s_{k,n}/n!$ is the degree *k* term of the Chern character of the tautological vector bundle E_n . Let us write

$$s_k = (s_{k,0}, s_{k,1}, \ldots) \in \prod_{n \ge 0} H^{2k}_{\mathscr{H}}(\mathrm{BGL}(n), \mathbb{R}(k))$$

Proposition 5.7. (Schechtman [Sch]) There exist elements $s_k^1 \in H_{\mathscr{H}}^{2k+1}$ $(\mathscr{B}P, \mathbb{R}(k))$ such that $d_H(s_k^1) = s_k$ and

$$H^*_{\mathscr{H}}(\mathscr{B}P, \mathbb{R}(*)) = A[s_0^1, s_1^1, \ldots]$$

(5.2.5) Since

$$H^n_{\mathscr{H}}(\operatorname{Spec}\mathbb{C},\mathbb{R}(p)) \cong \begin{cases} \mathbb{R}, & \text{if } n = p = 0, \text{ or } n = 1, p > 0, \\ 0, & \text{otherwise} \end{cases}$$

any element of $H^{2k+1}_{\mathscr{H}}(\mathscr{B}P,\mathbb{R}(k))$ can be written as

$$\alpha s_k^1$$
 + decomposable elements ,

with $\alpha \in \mathbb{R}$. Moreover, since by the proof of 5.7 [Sch] the decomposable elements are mapped to 0 by d_H , we have

Corollary 5.8. The group Ker $d_H \subset \bigoplus_k H^{2k+1}_{\mathscr{H}}(\mathscr{B}P, \mathbb{R}(k))$ is generated by decomposable elements.

(5.2.6) Schechtman computes the groups $\widehat{K}_*(\mathscr{B}P)$ in a similar way. In particular, there is also an edge homomorphism

$$d_K:\widehat{K}_{-1}(\mathscr{B}P)\longrightarrow\prod_n\widehat{K}_0(\mathrm{BGL}(n))$$

Moreover, by the naturality of ch and ρ , they are compatible with the edge homomorphisms. In particular

$$d_H(\rho(e_{\mathscr{B}P})) = \rho(d_K(e_{\mathscr{B}P})), \text{ and } d_H(\operatorname{ch}(e_{\mathscr{B}P})) = \operatorname{ch}(d_K(e_{\mathscr{B}P}))$$

(5.2.7) Our next step will be to compare $d_H(\rho(e_{\mathscr{B}P}))$ with $d_H(ch(e_{\mathscr{B}P}))$. To this end we shall see that, since the maps ch and ρ agree for the K_0 groups of smooth complex varieties then they also agree for the group $\widehat{K}_0(BGL(n))$.

Proposition 5.9. Let $\sigma \in \widehat{K}_0(BGL(n))$. Then

$$\operatorname{ch}(\sigma) = \rho(\sigma)$$
.

Proof. Let Gr(n,k) be the Grassman manifold of dimension *n* linear subspaces of C^k and let E(n,k) be the rank *n* tautological vector bundle. Let $\mathfrak{U}_k = \{U_\alpha\}$ be the standard trivialization of E(n,k). Let us denote by $\psi: N\mathfrak{U}_k \longrightarrow Gr(n,k)$ the natural map and by $\varphi_k: N\mathfrak{U}_k \longrightarrow BGL(n)$ the classifying map. Since absolute Hodge cohomology can be computed as the cohomology of a Zariski sheaf, the map

$$\psi^*: H^*_{\mathscr{H}}(\operatorname{Gr}(n,k),\mathbb{R}(*)) \longrightarrow H^*_{\mathscr{H}}(N\mathfrak{U}_k,\mathbb{R}(*))$$

is an isomorphism. Moreover, for each i_0 there is a number k_0 , such that, for all $k \ge k_0$ and all $i \le i_0$ the map

$$\varphi_k^*: H^i_{\mathscr{H}}(\mathrm{BGL}(n), \mathbb{R}(*)) \longrightarrow H^i_{\mathscr{H}}(N\mathfrak{U}_k, \mathbb{R}(*))$$

is an isomorphism. But for $\sigma \in \widehat{K}_0(\mathrm{BGL}(n))$ we have

$$\varphi_k^*(\mathrm{ch}(\sigma)) = \mathrm{ch}(\varphi_k^*(\sigma)) = \rho(\varphi_k^*(\sigma)) = \varphi_k^*\rho((\sigma)) \ .$$

Since this is true for all *k* we have $ch(\sigma) = \rho(\sigma)$.

Combining 5.8 and 5.9 we get:

Corollary 5.10. The element $ch(e_{\mathscr{B}P}) - \rho(e_{\mathscr{B}P})$ belongs to Ker d_H . Therefore it is a sum of decomposable elements.

(5.2.8) To exploit the fact that $ch(e_{\mathscr{B}P}) - \rho(e_{\mathscr{B}P})$ is a sum of decomposable elements, we shall give a description of how a class in $H^*_{\mathscr{H}}(\mathscr{B}P, \mathbb{R}(*))$ determines a map between K-theory and absolute Hodge cohomology.

For any smooth simplicial scheme over \mathbb{C} , X, and integers n, p, the complex

$$\mathscr{H}^*(X,n,p) = \tau_{\leq 0}\mathfrak{H}^*(X,p)[n]$$

is a negatively graded cohomological complex. Let $\mathscr{H}_*(X, n, p)$ be the associated homological complex. Let us denote by $\mathscr{K}(X, n, p)$ the simplicial group obtained by Dold-Puppe from $\mathscr{H}_*(X, n, p)$. Then, for $i \geq 0$,

 $\pi_i \mathscr{K}(X, n, p) = H^{n-i}_{\mathscr{H}}(X, \mathbb{R}(p)) \quad .$

Let us fix a smooth complex variety X, and \mathfrak{U} an open covering of X. Let us denote by φ the tautological map

 $\varphi: N\mathfrak{U} \times \underline{\mathrm{Hom}}(N\mathfrak{U}, \mathscr{B}P) \longrightarrow \mathscr{B}P$.

Given any class $x \in H^n_{\mathscr{H}}(\mathscr{B}P, \mathbb{R}(p))$, we have a class

$$\varphi^*(x) \in H^n_{\mathscr{H}}(N\mathfrak{U} \times \underline{\operatorname{Hom}}(N\mathfrak{U}, \mathscr{B}P), \mathbb{R}(p)) = \\ \operatorname{Hom}_{Ho}(\operatorname{Hom}(N\mathfrak{U}, \mathscr{B}P), \mathscr{K}(N\mathfrak{U}, n, p)) .$$

For any integer *i*, let us denote by $\pi_i(x)$ the induced map

$$\pi_i(x): K_{i-1}(X, \mathfrak{U}) = \pi_i \underline{\operatorname{Hom}}(N\mathfrak{U}, \mathscr{B}P) \longrightarrow \pi_i \mathscr{K}(N\mathfrak{U}, n, p) = H^{n-i}_{\mathscr{H}}(X, \mathbb{R}(p))$$

Taking the limit over all coverings we obtain morphisms

$$\pi_i(x): K_{i-1}(X) \longrightarrow H^{n-1}_{\mathscr{H}}(X, \mathbb{R}(p)) \quad .$$

This construction can be extended to the case when X is a simplicial smooth complex manifold.

Lemma 5.11. For $x \in H^{2k+1}_{\mathscr{H}}(\mathscr{B}P, \mathbb{R}(p))$ and $\sigma \in K_{i-1}(X, \mathfrak{U})$ we have $\pi_i(x)(\sigma) = \operatorname{pr}(\gamma^*_{\sigma}(x))$,

where γ_{σ} is as in (5.2.3).

Proof. Since the map $\pi_*(x)$ is natural, the same argument as for ch and ρ shows that

$$\pi_i(x)(\sigma) = \operatorname{pr}\big(\gamma^*_{\sigma}(\pi_0(x)(e_{\mathscr{B}P}))\big)$$

Let us denote by $\widehat{K}_i(\mathscr{BP}, \mathscr{BP})$ the K-theory groups of \mathscr{BP} with respect to the trivial covering. Then the map

$$\pi_0(x): \widehat{K}_{-1}(\mathscr{B}P, \mathscr{B}P) = \pi_0 \underline{\operatorname{Hom}}(\mathscr{B}P, \mathscr{B}P) \longrightarrow \pi_0 \mathscr{K}(\mathscr{B}P, 2k+1, k)$$
$$= H^{2k+1}(\mathscr{B}P, \mathbb{R}(k)) \quad ,$$

sends the class of $f \in \pi_0 \underline{\text{Hom}}(\mathscr{B}P, \mathscr{B}P)$ to $f^*(x)$. Since $e_{\mathscr{B}P}$ is represented by the identity map, we get

$$\pi_0(x)(e_{\mathscr{B}P}) = \mathrm{Id}^*(x) = x \; ,$$

proving the lemma.

(5.2.9) The product structure in absolute Hodge cohomology is given by a morphism of complexes

$$\mathscr{H}^*(X, n, p) \otimes \mathscr{H}^*(X, m, q) \xrightarrow{\cup} \mathscr{H}^*(X, n + m, p + q)$$
,

which induces a map of spaces

(1.)

$$\mathscr{K}^*(X,n,p) \times \mathscr{K}^*(X,m,q) \stackrel{\cup}{\longrightarrow} \mathscr{K}^*(X,n+m,p+q)$$

The spaces $\mathscr{K}(X, n, p)$ are naturally pointed by the element 0. Moreover $0 \cup x = x \cup 0 = 0$. Therefore the above map of spaces factors through:

$$\mathscr{H}^*(X,n,p) \times \mathscr{H}^*(X,m,q) \longrightarrow \mathscr{H}^*(X,n,p) \wedge \mathscr{H}^*(X,m,q) \longrightarrow \mathscr{H}^*(X,n+m,p+q) .$$

Lemma 5.12. Let $x \in H^n_{\mathscr{H}}(\mathscr{B}P, \mathbb{R}(p))$ and $y \in H^m_{\mathscr{H}}(\mathscr{B}P, \mathbb{R}(q))$. Then for any i > 0 the map $\pi_i(x \cup y) = 0$.

Proof. Let us write $E = \underline{\text{Hom}}(N\mathfrak{U}, \mathscr{B}P)$. Then the map $\pi(x \cup y)$ can be factored as

$$\pi_i(E) \xrightarrow{\pi_i(\operatorname{diag})} \pi_i(E \wedge E) \longrightarrow \pi_i(\mathscr{K}(N\mathfrak{U}, n, p) \wedge \mathscr{K}(N\mathfrak{U}, m, q)) \\ \longrightarrow \pi_i(\mathscr{K}(N\mathfrak{U}, n + m, p + q)) \quad .$$

292

But since $S^i \wedge S^i = S^{2i}$ and for i > 0, $\pi_i S^{2i} = 0$, the map $\pi_i(\text{diag}) = 0$.

(5.2.9) We are ready to prove Theorem 5.2. Let i > 0 and $\sigma \in K_{i-1}(X, \mathfrak{U})$. By Lemma 5.11, we have that

$$\mathrm{ch}(\sigma) = \pi_i(\mathrm{ch}(e_{\mathscr{B}P}))(\sigma), \
ho(\sigma) = \pi_i(
ho(e_{\mathscr{B}P}))(\sigma) \; .$$

Therefore

$$\operatorname{ch}(\sigma) - \rho(\sigma) = \pi_i(\operatorname{ch}(e_{\mathscr{B}P}) - \rho(e_{\mathscr{B}P}))(\sigma)$$

By Corollary 5.10, $ch(e_{\mathscr{B}P}) - \rho(e_{\mathscr{B}P})$ is a sum of decomposable elements. Therefore by Lemma 5.12.

$$ch(\sigma) = \rho(\sigma)$$

concluding the proof of the theorem.

(5.3) The same argument shows that, for a smooth simplicial complex variety X, an integer $i \ge 0$, and an element $\sigma \in \widehat{K}_i(X)$ then $ch(\sigma) = \rho(\sigma)$. To prove the same result for i < 0 one can use an analogous argument using $\mathscr{B}^m P$ ([Sch]), with m > -i.

6 Higher Bott-Chern forms

The higher Bott-Chern forms introduced in Sect. 3 are differential forms defined on $X \times (\mathbb{P}^1)^*$. Nevertheless, the original Bott-Chern forms [B-C] and the higher Bott-Chern forms introduced by Wang in [Wan] are differential forms defined on X. The aim of this section is to relate both notions of higher Bott-Chern forms, in the case when X is a proper smooth complex variety. The main tool for this comparison will be an explicit quasi-isomorphism

$$\mathfrak{H}^*(X,p) \longrightarrow \mathfrak{H}^*(X,p)$$
.

To this end we shall first introduce some complexes which compute absolute Hodge homology and cohomology.

(6.1) Let us begin by introducing the complex where the simplest Bott-Chern forms are defined. This complex is a minor modification of the complex used by Wang in [Wan] (see also [Bu 2]). The use of this complex has been suggested by Deligne in [De]. Let X be a proper smooth complex variety. We shall write

$$E^*_{\mathbb{R}}(X)(p) = (2\pi i)^p E^*_{\mathbb{R}}(X)$$
.

Definition 6.1. The complex $\mathfrak{W}^*(X,p)$ is defined by

$$\mathfrak{W}^{n}(X,p) = \begin{cases} E_{\mathbb{R}}^{n-1}(X)(p-1) \cap \bigoplus_{\substack{p'+q'=n-1\\p' < p, \ q' < p}} E_{p',q'(X)}^{p',q'(X)}, & \text{for } n \leq 2p-1, \\ E_{\mathbb{R}}^{n}(X)(p) \cap \bigoplus_{\substack{p'+q'=n\\p' \geq p, \ q' \geq p}} E_{p',q'(X)}^{p',q'(X)} \cap \operatorname{Ker} d, & \text{for } n = 2p, \\ 0, & \text{for } n > 2p . \end{cases}$$

If $x \in \mathfrak{W}^n(X,p)$ the differential $d_{\mathfrak{W}}$ is given by

$$d_{\mathfrak{M}}x = \begin{cases} -\pi(dx), & \text{for } n < 2p - 1, \\ -2\partial\overline{\partial}x, & \text{for } n = 2p - 1, \\ 0, & \text{for } n = 2p \end{cases},$$

where

$$\pi: E^*(X) \longrightarrow E^*_{\mathbb{R}}(X)(p-1) \cap \bigoplus_{\substack{p'+q'=n-1\\p' < p, \ q' < p}} E^{p',q'}(X) \ ,$$

is the projection.

Proposition 6.2. If X is a proper smooth complex variety, then

$$H^*(\mathfrak{W}^*(X,p)) = H^*_{\mathscr{H}}(X,\mathbb{R}(p)) \ .$$

Proof. Since X is proper,

$$H^n_{\mathscr{H}}(X, \mathbb{R}(p)) = \begin{cases} H^n_{\mathscr{D}}(X, \mathbb{R}(p)), & \text{for } n \le 2p, \\ 0, & \text{for } n > 2p, \end{cases}$$

where $H^n_{\mathscr{D}}(X, \mathbb{R}(p))$, denotes real Deligne cohomology of X. Therefore the result follows from [Bu 2, Sect. 2].

As in [Bu 2], we have morphisms of complexes

$$\psi: \mathfrak{H}^*(X,p) \longrightarrow \mathfrak{W}^*(X,p)$$

and

$$\varphi: \mathfrak{W}^*(X,p) \longrightarrow \mathfrak{H}^*(X,p)$$

given by

$$\psi(a, f, \omega) = \begin{cases} \pi(\omega), & \text{for } n \le 2p - 1\\ & \text{and} \\ \sum_{i=p}^{n-p} a^{i,n-i} + \partial \omega^{p-1,n-p+1} + (-1)^p \overline{\partial} \overline{\omega}^{p-1,n-p+1}, & \text{for } n \ge 2p \end{cases},$$

294

and

$$\varphi(x) = \begin{cases} \left(\partial x^{p-1,n-p} - \overline{\partial} x^{n-p,p-1}, 2\partial x^{p-1,n-p}, x\right), & \text{for } n \le 2p-1 \text{ and} \\ (x,x,0), & \text{for } n \ge 2p \end{cases}$$

where, if $x \in E_X^*$, then $x = \sum x^{p,q}$ is the decomposition of x in terms of pure type. The morphisms φ and ψ are homotopy equivalences inverse to each other.

(6.2) In order to make the process of comparison clearer, we need an auxiliary complex to compute absolute Hodge cohomology, which is provided with a graded commutative and associative product. It can be obtained by means of the Thom-Whitney simple introduced by Navarro Aznar (see [N 2] for the general definition and properties of the Thom-Whitney simple).

Let L_1^* be the differential graded commutative \mathbb{R} -algebra of algebraic forms over $\mathbb{A}^1_{\mathbb{R}}$. Explicitly $L_1^0 = \mathbb{R}[\epsilon]$ and $L_1^1 = \mathbb{R}[\epsilon]d\epsilon$. Let $\delta_0 : L_1^* \longrightarrow \mathbb{R}$ (resp. δ_1) be the evaluation at 0 morphism (resp. evaluation at 1).

Definition 6.3. Let X be a smooth complex variety. The Thom-Whitney simple of the absolute Hodge complex, denoted by $\mathfrak{H}^*_{TW}(X,p)$, is the subcomplex of

$$\left((2\pi i)^p \widehat{W}_{2p} E^*_{\log}(X)_{\mathbb{R}} \oplus \widehat{W}_{2p} \cap F^p E^*_{\log}(X) \oplus \left(L^*_1 \underset{\mathbb{R}}{\otimes} \widehat{W}_{2p} E^*_{\log}(X) \right) \right)$$

formed by the elements (r, f, ω) such that

$$\omega(0) = (\delta_0 \otimes \mathrm{Id})(\omega) = r,$$

$$\omega(1) = (\delta_1 \otimes \mathrm{Id})(\omega) = f .$$

Let E and I be the morphisms of complexes

$$\mathfrak{H}^*_{\mathrm{TW}}(X,p) \xrightarrow[E]{\mathrm{I}} \mathfrak{H}^*(X,p)$$

given by

$$\begin{split} E(r,f,\omega) &= (r,f,\epsilon\otimes f + (1-\epsilon)\otimes r + d\epsilon\otimes \omega),\\ I(r,f,\omega) &= \left(r,f,\int_{0}^{1}\omega\right) \;, \end{split}$$

where the integration symbol means formal integration with respect to the variable ϵ . These morphisms are homotopy equivalences (see [N 2]).

We shall denote by I' the composition

$$\mathfrak{H}^*_{\mathrm{TW}}(X,*) \xrightarrow{I} \mathfrak{H}^*(X,*) \xrightarrow{\psi} \mathfrak{W}^*(X,*)$$

and by E' the composition

$$\mathfrak{W}^*(X,*) \xrightarrow{\varphi} \mathfrak{H}^*(X,*) \xrightarrow{E} \mathfrak{H}^*_{\mathrm{TW}}(X,*)$$
.

The morphisms I' and E' are also homotopy equivalences inverse to each other.

We can define a product

$$\mathfrak{H}^n_{\mathrm{TW}}(X,p)\otimes\mathfrak{H}^m_{\mathrm{TW}}(X,q) \xrightarrow{\cup} \mathfrak{H}^{n+m}_{\mathrm{TW}}(X,p+q) \;,$$

by

$$(r, f, \omega) \cup (r', f', \omega') = (r \wedge r', f \wedge f', \omega \wedge \omega')$$

This product is associative, graded commutative and satisfies the Leibnitz rule. Therefore

$$\mathfrak{H}^*_{\mathrm{TW}}(X,*) = \bigoplus_p \mathfrak{H}^*_{\mathrm{TW}}(X,p)$$

is a differential associative graded commutative algebra. Moreover, the \mathbb{R} -algebra structure induced in $H^*_{\mathscr{H}}(X, \mathbb{R}(p))$ by this product coincides with the \mathbb{R} -algebra structure introduced by Beilinson [Be].

(6.3) Let us give the homology analogue of the last complex. This is done by means of currents. For a proper smooth complex variety X, let $D_{*,*}(X)$ be the double chain complex of complex currents over X, let $D_*(X)$ be the associated single complex, and let $D_*^{\mathbb{R}}(X)$ be the real subcomplex. We shall write

$$F_p D_*(X) = \bigoplus_{p' \le p} D_{p',*}$$
.

Let $\tau^{\geq 2p}D_*(X)$ be the subcomplex

$$\tau^{\geq 2p} D_n(X) = \begin{cases} D_n(X), & \text{if } n > 2p, \\ \text{Ker}(d), & \text{if } n = 2p, \\ 0, & \text{if } n < 2p \end{cases}$$

Since X is proper, the filtration τ plays the role of the décalée weight filtration.

Let L_*^1 be the chain complex defined by $L_k^1 = L_1^{-k}$ (see 6.2). We shall denote by δ_0 and δ_1 the evaluation at 0 and 1 as in (6.2).

Definition 6.4. Let $\mathfrak{H}^{\mathrm{TW}}_*(X,p)$ be the subcomplex of

$$\left((2\pi i)^{-p}\tau^{\geq 2p}D^{\mathbb{R}}_{*}(X)\oplus\tau^{\geq 2p}\cap F_{p}D_{*}(X)\oplus\left(L^{1}_{*}\otimes\tau^{\geq 2p}D_{*}(X)\right)\right)$$

formed by the elements (r, f, ω) such that

$$\omega(0) = (\delta_0 \otimes \mathrm{Id})(\omega) = r,$$

$$\omega(1) = (\delta_1 \otimes \mathrm{Id})\omega = f .$$

The homology of the complex $\mathfrak{H}^{\mathrm{TW}}_*(X,p)$ is the absolute Hodge homology of X.

(6.4) The last complex we introduce is an analogue of $\tilde{\mathfrak{H}}^*(X,p)$, replacing $\mathfrak{H}^*(X,p)$ by $\mathfrak{H}^*_{TW}(X,p)$. We shall denote by $\mathfrak{H}^{**}_{\mathbb{P},TW}(X,p)$ the double complex given by

$$\mathfrak{H}^{r,n}_{\mathbb{P},\mathrm{TW}}(X,p) = \mathfrak{H}^{r}_{\mathrm{TW}}(X \times (\mathbb{P}^{1})^{-n},p)$$

with differentials

$$d' = d_{\mathfrak{H}}, \ d'' = \sum (-1)^{i+j} d_i^j \;\;.$$

Then the double complex $\widetilde{\mathfrak{H}}_{\mathrm{TW}}^{*,*}(X,p)$ is given by

$$\widetilde{\mathfrak{H}}_{\mathsf{TW}}^{r,n}(X,p) = \mathfrak{H}_{\mathbb{P},\mathsf{TW}}^{r,n}(X,p) \left/ \sum_{i=1}^{n} s_i \Big(\mathfrak{H}_{\mathbb{P},\mathsf{TW}}^{r,n+1}(X,p) \Big) \oplus \omega_i \wedge s_i \Big(\mathfrak{H}_{\mathbb{P},\mathsf{TW}}^{r-2,n+1}(X,p-1) \Big) \right.$$

Finaly let $\tilde{\mathfrak{F}}_{TW}^*(X,p)$ be the associated simple complex. The differential of this complex will be denoted by *d*.

Observe that the homotopy equivalences E and I induce homotopy equivalences

$$\begin{array}{c} \overset{\mathrm{I}}{\mathfrak{H}}_{\mathrm{TW}}^{*}(X,p) \xrightarrow[]{E}]{} \overset{\mathrm{I}}{\underset{\mathrm{E}}{\longrightarrow}} \widetilde{\mathfrak{H}}^{*}(X,p) \ . \end{array}$$

(6.5) In order to pull down forms in $X \times (\mathbb{P}^1)^n$ to X, we need some differential forms on $X \times (\mathbb{P}^1)^n$ which will play a role similar to the currents "integration along the standard simplex".

Let (x : y) be homogeneous coordinates of \mathbb{P}^1 , and let t = x/y be the absolute coordinate of \mathbb{P}^1 . Let us write $\mathbb{C}^* = \mathbb{P}^1_{\mathbb{C}} - \{0, \infty\}$. Let

$$\begin{split} \lambda &= \frac{1}{2} E'(\log t\overline{t}) \\ &= \frac{1}{2} \left(\frac{dt}{t} - \frac{d\overline{t}}{\overline{t}}, 2\frac{dt}{t}, (\varepsilon + 1) \otimes \frac{dt}{t} + (\varepsilon - 1) \otimes \frac{d\overline{t}}{\overline{t}} + d\varepsilon \otimes \log t\overline{t} \right) \\ &\in \mathfrak{H}_{\mathrm{TW}}^{1}(\mathbb{C}^{*}, 1) \quad . \end{split}$$

Let us consider the open subset $(\mathbb{C}^*)^n \subset X \times (\mathbb{P}^1)^n$. Let us denote by $p_i : (\mathbb{C}^*)^n \longrightarrow \mathbb{C}^*$, i = 1, ..., n the projections over the *i*-th factor. Let us write $\lambda_i = p_i^* \lambda$.

Definition 6.5. Let $W_n \in \mathfrak{H}^n_{TW}((\mathbb{C}^*)^n, n)$ be the form defined by

$$W_n = \lambda_1 \cup \cdots \cup \lambda_n$$
.

(6.6) Since the forms W_n will play a central role, let us present a more explicit description. Let us write $W_n = (W_n^1, W_n^2, W_n^3)$. Then

$$W_n^1 = \frac{1}{2^n} \bigwedge_{i=1}^n \left(\frac{dt_i}{t_i} - \frac{d\overline{t_i}}{\overline{t_i}} \right)$$
$$W_n^2 = \bigwedge_{i=1}^n \frac{dt_i}{t_i}$$
$$W_n^3 = \frac{1}{2^n} \bigwedge_{1=1}^n \left((\epsilon + 1) \otimes \frac{dt_i}{t_i} + (\epsilon - 1) \otimes \frac{d\overline{t_i}}{\overline{t_i}} + d\epsilon \otimes \log t_i \overline{t_i} \right)$$

Let \mathfrak{S}_n denote the symmetric group. Let us write, for $i = 0, \ldots, n$,

$$P_n^i = \sum_{\sigma \in \mathfrak{S}_n} (-1)^{\sigma} \frac{dt_{\sigma(1)}}{t_{\sigma(1)}} \wedge \cdots \wedge \frac{dt_{\sigma(i)}}{t_{\sigma(i)}} \wedge \frac{d\overline{t}_{\sigma(i+1)}}{\overline{t}_{\sigma(i+1)}} \wedge \cdots \wedge \frac{d\overline{t}_{\sigma(n)}}{\overline{t}_{\sigma(n)}} ,$$

and, for i = 1, ..., n,

$$S_n^i = \sum_{\sigma \in \mathfrak{S}_n} (-1)^{\sigma} \log \left(t_{\sigma(1)} \overline{t}_{\sigma(1)} \right) \frac{dt_{\sigma(2)}}{t_{\sigma(2)}} \wedge \dots \wedge \frac{dt_{\sigma(i)}}{t_{\sigma(i)}} \wedge \frac{d\overline{t}_{\sigma(i+1)}}{\overline{t}_{\sigma(i+1)}} \wedge \dots \wedge \frac{d\overline{t}_{\sigma(n)}}{\overline{t}_{\sigma(n)}}$$

Then we have

$$\begin{split} W_n^1 &= \frac{1}{2^n} \sum_{i=0}^n (-1)^{n-i} \frac{1}{i!(n-i)!} P_n^i, \\ W_n^3 &= \frac{1}{2^n} \sum_{i=0}^n \frac{(\epsilon+1)^i (\epsilon-1)^{n-i}}{i!(n-i)!} \otimes P_n^i + \frac{1}{2^n} \sum_{i=1}^n \frac{(\epsilon+1)^{i-1} (\epsilon-1)^{n-i}}{(i-1)!(n-i)!} d\epsilon \otimes S_n^i \end{split}$$

(6.7) We are not as interested in the forms W_n , as in their associated currents. Let $\omega \in E^r_{(\mathbb{P}^1)^n}$. Let us denote by $[\omega] \in D_{2n-r}((\mathbb{P}^1)^n)$ the current defined by

298

$$[\omega](\varphi) = \frac{1}{(2\pi i)^n} \int_{(\mathbb{P}^1)^n} \varphi \wedge \omega \ .$$

If $a \otimes \omega \in L_1^* \otimes E_{(\mathbb{P}^1)^n}^r$ we write

$$egin{aligned} [a\otimes\omega]&=a\otimes[\omega]\ &\in L^1_*\otimes D_{2n-r}ig(ig({f P}^1ig)^nig) \end{aligned}$$

In this way we obtain a map

$$\mathfrak{H}^{r}_{\mathrm{TW}}((\mathbb{P}^{1})^{n}, p) \longrightarrow \mathfrak{H}^{\mathrm{TW}}_{2n-r}((\mathbb{P}^{1})^{n}, n-p) \\
(r, f, \omega) \longmapsto ([r], [f], [\omega])$$

This definition can be extended to any locally integrable differential form.

Definition 6.6. We shall denote by $[W_n]$ the element of $\mathfrak{H}_n^{\mathrm{TW}}((\mathbb{P}^1)^n, 0)$ given by

$$[W_n] = \left(\begin{bmatrix} W_n^1 \end{bmatrix}, \begin{bmatrix} W_n^2 \end{bmatrix}, \begin{bmatrix} W_n^3 \end{bmatrix} \right) .$$

The following result exhibits the analogy between the currents "integration along the standard simplex" and the currents $[W_n]$.

Proposition 6.7. The currents $[W_n]$ satisfy the relation

$$d[W_n] = \sum_{i=1}^n \sum_{j=0,1} (-1)^{i+j} \left(d_j^i \right)_* [W_{n-1}] .$$

Proof. Formally this proposition is the Leibnitz rule. To prove it we can work component by component. By a standard residue argument:

$$d[W_n^2] = d\left[\bigwedge_{i=1}^n \frac{dt_i}{t_i}\right]$$
$$= \sum_{i=1}^n \sum_{j=0}^{1} (-1)^{i+j} \left(d_j^i\right)_* [W_{n-1}^2]$$

By the same argument and taking some care with permutations one sees

$$d[P_n^i] = \sum_{k=1}^n \sum_{j=0}^{1} (-1)^{k+j} \left(d_j^k \right)_* \left(i [P_{n-1}^{i-1}] - (n-i) [P_{n-1}^i] \right),$$

$$d[S_n^i] = [P_n^i] + [P_n^{i-1}] + \sum_{k=1}^n \sum_{j=0}^{1} (-1)^{k+j} \left(d_j^k \right)_* \left((i-1) [S_{n-1}^{i-1}] - (n-i) [S_{n-1}^i] \right).$$

The proposition follows from the above formulas and the explicit description of W_n^1 and W_n^3 given in (6.6).

(6.8) Acting component by component, the currents $[W_n]$ induce morphisms

$$[W_n]: \mathfrak{H}^{r,-n}_{\mathbb{P},\mathrm{TW}}(X,p) = \mathfrak{H}^r_{\mathrm{TW}}(X \times (\mathbb{P}^1)^n,p) \longrightarrow \mathfrak{H}^{r-n}_{\mathrm{TW}}(X,p)$$

Lemma 6.8. The morphisms $[W_n]$ factorize through morphisms

$$[W_n]: \mathfrak{H}^{r,n}_{\mathrm{TW}}(X,p) \longrightarrow \mathfrak{H}^{r-n}_{\mathrm{TW}}(X,p)$$

Proof. Let us denote by σ_i the automorphism of $(\mathbb{P}^1)^n$ given by

$$\sigma_i((x_1:y_1),\ldots,(x_i:y_i),\ldots,(x_n:y_n)) = ((x_1:y_1),\ldots,(y_i:x_i),\ldots,(x_n:y_n))$$

Then $(\sigma_i)_*[W_n] = -[W_n]$, for i = 1, ..., n. On the other hand, if

$$\eta \in s_i\left(\mathfrak{H}^{r,n+1}_{\mathbb{P},\mathrm{TW}}(X,p)\right) \oplus \omega_i \wedge s_i\left(\mathfrak{H}^{r-2,n+1}_{\mathbb{P},\mathrm{TW}}(X,p-1)\right)$$

then $(\sigma_i)^*\eta = \eta$. Therefore

$$[W_n]\eta = -(\sigma_i)_*[W_n]\eta = -[W_n](\sigma_i)^*\eta = -[W_n]\eta$$
.

Hence $[W_n]\eta = 0$ proving the result.

Definition 6.9. Let W_{TW} be the morphism

$$W_{\mathrm{TW}}: \widetilde{\mathfrak{H}}^*_{\mathrm{TW}}(X,p) \longrightarrow \mathfrak{H}^*_{\mathrm{TW}}(X,p)$$

given, for $\eta \in \widetilde{\mathfrak{H}}_{\mathrm{TW}}^{r,n}(X,p)$, by

$$W_{\mathrm{TW}}(\eta) = egin{cases} [W_n]\eta, & ext{if } n > 0, \ \eta, & ext{if } n = 0 \end{cases}$$

Proposition 6.10. The morphism W_{TW} is a morphism of complexes. Moreover it is a quasi-isomorphism.

Proof. The fact that is a morphism of complexes is a consequence of Proposition 6.7. Let

$$\iota':\mathfrak{H}^*_{\mathrm{TW}}(X,p)\longrightarrow \widetilde{\mathfrak{H}}^*_{\mathrm{TW}}(X,p)$$

be the morphism induced by the equality $\mathfrak{H}^{*,0}_{\mathrm{TW}}(X,p) \longrightarrow \widetilde{\mathfrak{H}}^{*,0}_{\mathrm{TW}}(X,p)$. We have that $\iota = I \circ \iota' \circ E$, where ι is the quasi-imorphism defined in Proposition 1.2.

Therefore $\iota' = \text{Id}$ is a quasi-isomorphism. Since $W_{\text{TW}} \circ \iota' = \text{Id}$ we have that W_{TW} is also a quasi-isomorphism.

Definition. 6.11. Let us denote by W the morphism

$$W = I \circ W_{\mathrm{TW}} \circ E : \widetilde{\mathfrak{H}}^*(X, p) \longrightarrow \mathfrak{H}^*(X, p) \ .$$

Observe that W is also a quasi-isomorphism. Summarizing, we have the following diagram of complexes and quasi-isomorphisms.

$$\begin{array}{ccccc} \widetilde{\mathfrak{H}}^{*}(X,p) & \xrightarrow{W} & \mathfrak{H}^{*}(X,p) & \xrightarrow{\psi} & \mathfrak{W}^{*}(X,p) \\ E & & I \\ \widetilde{\mathfrak{H}}^{*}_{\mathrm{TW}}(X,p) & \xrightarrow{W_{\mathrm{TW}}} & \mathfrak{H}^{*}_{\mathrm{TW}}(X,p) \end{array}$$

(6.9) The diagram above allows us to define different kinds of higher Bott-Chern forms. For instance let us recover the original definition of higher Bott-Chern forms due to Wang [Wan] and the classical Bott-Chern forms.

Definition 6.12. Let $\overline{\mathcal{F}}$ be an exact metrized n-cube. We shall also call the Bott-Chern form of $\overline{\mathcal{F}}$ the form

$$\widetilde{\mathrm{ch}}(\overline{\mathscr{F}})_W = \psi \circ W\left(\widetilde{\mathrm{ch}}_n(\overline{\mathscr{F}})_{\mathscr{H}}\right)$$
.

One may compute these forms directly using the following result.

Proposition 6.13. Let $\overline{\mathcal{F}}$ be an emi-*n*-cube. Then

$$\widetilde{\mathrm{ch}}_n(\overline{\mathscr{F}})_W = \frac{1}{(2\pi i)^n} \int_{\left(\mathbb{P}^1\right)^n} \widetilde{\mathrm{ch}}_0(\mathrm{tr}_n(\overline{\mathscr{F}})) \wedge I'(W_n) .$$

Proof. This result is consequence of the following facts

(1) The morphism I' is functorial.

(2) For any smooth complex variety Z, if $\omega \in \mathfrak{H}^{2p}_{\mathrm{TW}}(Z,p)$ and $\eta \in \mathfrak{H}^*_{\mathrm{TW}}(Z^*,*)$, then $I'(\omega \cup \eta) = I'(\omega) \wedge I'(\eta)$ (see [Bu 2]).

(3) $I' \circ E' =$ Id. Therefore

$$I'(E(\widetilde{ch}_0(\mathrm{tr}_n(\overline{\mathscr{F}}))_{\mathscr{H}})) = I'(E'(\widetilde{ch}_0(\mathrm{tr}_n(\overline{\mathscr{F}})))) = \widetilde{ch}_0(\mathrm{tr}_n(\overline{\mathscr{F}})) \ .$$

Up to a normalization factor, the formula given in Proposition 6.13 is the original definition due to Wang [Wan]. To see this, let us compute explicitly $I'(W_n) \in \mathfrak{M}^n((\mathbb{C}^*)^n, n)$.

Proposition 6.14.

$$I'(W_n) = \frac{(-1)^n}{2n!} \sum_{i=1}^n (-1)^{i-1} S_n^i \ .$$

Proof. Since $W_n \in \mathfrak{H}^n_{\mathrm{TW}}((\mathbb{C}^*)^n, n)$, by (6.1) and (6.2), we have

$$I'(W_n) = \pi \left(\int_0^1 W_n^3 \right) \; ,$$

where the integral symbol means integration with respect to the variable ϵ , and π is the projection

$$\pi: E^{n-1}_{(\mathbb{C}^*)^n} \longrightarrow (2\pi i)^{n-1} E^{n-1}_{(\mathbb{C}^*)^n, \mathbb{R}}$$

This projection is given by $\pi(z) = (z + (-1)^{n-1}\overline{z})/2$. Therefore

$$I'(W_n) = \frac{1}{2^{n+1}} \sum_{i=1}^n \int_0^1 \frac{(\epsilon+1)^{i-1}(\epsilon-1)^{n-i}}{(i-1)!(n-i)!} d\epsilon \left(S_n^i + (-1)^{n-1}\overline{S}_n^i\right) .$$

But $\overline{S}_n^i = S_n^{n-i+1}$. Then, joining the terms with S_n^i , and taking into account that

$$(-1)^{n-1} \int_0^1 \frac{(\epsilon+1)^{n-i}(\epsilon-1)^{i-1}}{(i-1)!(n-i)!} d\epsilon = \int_{-1}^0 \frac{(\epsilon+1)^{i-1}(\epsilon-1)^{n-i}}{(i-1)!(n-i)!} d\epsilon ,$$

we have that

$$I'(W_n) = \frac{1}{2^{n+1}} \sum_{i=1}^n \left(\int_{-1}^1 \frac{(\epsilon+1)^{i-1}(\epsilon-1)^{n-i}}{(i-1)!(n-i)!} \ d\epsilon S_n^i \right) \ .$$

But

$$\int_{-1}^{1} \frac{(\epsilon+1)^{i-1}(\epsilon-1)^{n-i}}{(i-1)!(n-i)!} \ d\epsilon = \frac{(-1)^{n+i-1}2^n}{n!} \ ,$$

proving the result.

The following result is a direct consequence of the definitions.

Proposition 6.15. Let X be a proper smooth complex variety. Let

 $\overline{\xi}: 0 \longrightarrow \overline{\mathscr{F}} \longrightarrow \overline{\mathscr{G}} \longrightarrow \overline{\mathscr{H}} \longrightarrow 0$

be an exact sequence of locally free sheaves over X. Let us denote by $\widetilde{bc}(\overline{\xi})$ the Bott-Chern form of $\overline{\xi}$ as defined by Bismut, Gillet and Soulé [B-G-S], [G-S 1]). Then

302

$$\widetilde{\mathrm{ch}}_1(\overline{\xi})_W = -1/2 \ \widetilde{\mathrm{bc}}\overline{\xi} \qquad \mathrm{mod}(\mathrm{Im}\,\partial + \mathrm{Im}\,\overline{\partial}) \ .$$

(6.10) The use of the Thom-Whitney simple for absolute Hodge cohomology, besides giving a way to construct the currents W_n , allows us to define a multiplicative theory of Bott-Chern forms.

Definition 6.16. Let $\overline{\mathcal{F}}$ be an exact metrized n-cube. We shall call the multiplicative Bott-Chern form of $\overline{\mathcal{F}}$ the form

$$\widetilde{\mathrm{ch}}_n(\overline{\mathscr{F}})_{\mathrm{TW}} = W_{\mathrm{TW}}(E(\widetilde{\mathrm{ch}}_n(\overline{\mathscr{F}})_{\mathscr{H}}))$$
.

In particular, if $\overline{\mathcal{F}}$ is a hermitian locally free sheaf, then

$$\widetilde{\mathrm{ch}}_0(\overline{\mathscr{F}})_{\mathrm{TW}} = E(\widetilde{\mathrm{ch}}_0(\overline{\mathscr{F}})_{\mathscr{H}})$$
.

On the other hand, if $\overline{\mathcal{F}}$ is an emi-n-cube, then

$$\widetilde{\mathrm{ch}}_n(\overline{\mathscr{F}})_{\mathrm{TW}} = \frac{1}{(2\pi i)^n} \int_{(\mathbb{P}^1)^n} \widetilde{\mathrm{ch}}_0(\mathrm{tr}_n(\overline{\mathscr{F}}))_{\mathrm{TW}} \cup W_n$$
.

Definition 6.17. Let $\overline{\mathcal{F}}$ be a metrized exact *n*-cube and let $\overline{\mathcal{G}}$ be a metrized exact *m*-cube. Then $\overline{\mathcal{F}} \otimes \overline{\mathcal{G}}$ is the metrized exact n + m-cube given by

$$(\overline{\mathscr{F}}\otimes\overline{\mathscr{G}})_{i_1,\ldots,i_{n+m}}=(\overline{\mathscr{F}})_{i_1,\ldots,i_n}\otimes(\overline{\mathscr{G}})_{i_{n+1},\ldots,i_{n+m}}$$

with the obvious morphisms and metrics.

Proposition 6.18. Let $\overline{\mathcal{F}}$ (resp. $\overline{\mathcal{G}}$) be a metrized exact n-cube (resp. m-cube). Then

$$\widetilde{ch}_{n+m}(\overline{\mathscr{F}}\otimes\overline{\mathscr{G}})_{\mathrm{TW}}=\widetilde{ch}_n(\overline{\mathscr{F}})_{\mathrm{TW}}\cup\widetilde{ch}_m(\overline{\mathscr{G}})_{\mathrm{TW}}$$

Proof. We may assume that $\overline{\mathscr{F}}$ and $\overline{\mathscr{G}}$ are emi-cubes.

Let $\pi_1: X \times (\mathbb{P}^1)^{n+m} \longrightarrow X \times (\mathbb{P}^1)^n$ be the projection over the first *n*-projective lines and let $\pi_2: X \times (\mathbb{P}^1)^{n+m} \longrightarrow X \times (\mathbb{P}^1)^m$ be the projection over the last *m*-projective lines.

Lemma 6.19. Let $\overline{\mathscr{F}}$ (resp. $\overline{\mathscr{G}}$) be an emi-*n*-cube (resp. emi-*m*-cube). Then

$$\mathrm{tr}_{n+m}(\overline{\mathscr{F}}\otimes\overline{\mathscr{G}})=\pi_1^*\mathrm{tr}_n(\overline{\mathscr{F}})\otimes\pi_2^*\mathrm{tr}_m(\overline{\mathscr{G}})$$

Proof. By Sect. 3, (3.7), it is enough to show that, if $m \ge 1$, then

$$\operatorname{tr}_1(\overline{\mathscr{F}}\otimes\overline{\mathscr{G}})=\overline{\mathscr{F}}\otimes\operatorname{tr}_1(\overline{\mathscr{G}})$$

and if m = 0, then

$$\operatorname{tr}_1(\overline{\mathscr{F}}\otimes\overline{\mathscr{G}}) = \operatorname{tr}_1(\overline{\mathscr{F}})\otimes\overline{\mathscr{G}}$$

Since tr_1 is computed in each edge separately, it is enough to prove the case n = 1, m = 0, but this case follows directly from the definition.

Using lemma 6.19, the multiplicativity and functoriality of the Chern character forms and the definition of the forms W_n , we have:

$$\begin{split} \widetilde{\mathrm{ch}}_{n+m}(\overline{\mathscr{F}}\otimes\overline{\mathscr{G}})_{\mathrm{TW}} &= \frac{1}{(2\pi i)^{n+m}} \int_{(\mathbb{P}^1)^{n+m}} \widetilde{\mathrm{ch}}_0(\pi_1^*\mathrm{tr}_n(\overline{\mathscr{F}})\otimes\pi_2^*\mathrm{tr}_m(\overline{\mathscr{G}}))_{\mathrm{TW}} \cup W_{n+m} \\ &= \frac{1}{(2\pi i)^{n+m}} \int_{(\mathbb{P}^1)^{n+m}} \pi_1^*\widetilde{\mathrm{ch}}_0(\mathrm{tr}_n(\overline{\mathscr{F}}))_{\mathrm{TW}} \cup \pi_2^*\widetilde{\mathrm{ch}}_0(\mathrm{tr}_m(\overline{\mathscr{G}}))_{\mathrm{TW}} \cup \pi_1^*W_n \cup \pi_2^*W_m \\ &= \widetilde{\mathrm{ch}}_n(\overline{\mathscr{F}})_{\mathrm{TW}} \cup \widetilde{\mathrm{ch}}_m(\overline{\mathscr{G}})_{\mathrm{TW}} \quad . \end{split}$$

References

- [Be] Beilinson, A.A.: Notes on absolute Hodge cohomology, Contemporary Mathematics 55 I, 35–68 (1968)
- [B-G-S] Bismut, J.M., Gillet, H., Soulé, C.: Analytic torsion and holomorphic determinant bundles I: Bott-Chern forms and analytic torsion, Comm. Math. Phys. 115, 49–78 (1988)
- [B-C] Bott, R., Chern, S.S.: Hermitian vector bundles and the equidistribution of zeroes of their holomorphic sections, Acta Math 114, 71–112 (1968)
- [B-K] Bousfield A.K., Kan D.M.: Homotopy limits, completions and localizations. LNM 304, Springer-Verlag, 1972
- [Bu 1] Burgos J.I.: A C[∞] logarithmic Dolbeault complex, Compos. Math. 92, 61–86 (1994)
- [Bu 2] Burgos J.I.: Arithmetic Chow rings and Deligne Beilinson cohomology, J. of Algebraic Geometry 6, 335–377 (1997)
- [Ch] Chern S.S.: Characteristic classes of hermitian manifolds, Ann. of Math. 47, 85–121 (1946)
- [De] Deligne P.: Le déterminant de la cohomologie, Contemporary Mathematics 67, 93– 177 (1987)
- [Gi] Gillet, H.: Riemann-Roch theorems for higher algebraic K-theory, Advances in Math. 40, 203–289 (1981)
- [G-S 1] Gillet, H., Soulé, C.: Characteristic classes for algebraic vector bundles with hermitian metric I and II, Annals of Math. 131, 163–238 (1990)
- [G-S 2] Gillet, H., Soulé, C.: Arithmetic intersection theory, Publ. Math. IHES 72, 93–174 (1990)
- [Gr] Griffiths, P.A.: On a theorem of Chern, Illinois J. Math. 6, 468–479 (1962)
- [G-D] Grothendieck, A., Dieudonné, J.A.: Eléments de géométrie algébrique, Springer-Verlag, 1971
- [GNPP] Guillen, F., Navarro Aznar, V., Pascual-Gainza, P., Puerta, F.: Hyperrésolutions cubiques et descente cohomologique. LNM 1335, Springer-Verlag, 1988
- [Ka 1] Karoubi, M.: Homologie cyclique et K-théorie Astérisque 149, 1987

- [Ka 2] Karoubi,: Théorie générale des classes caractéristiques secondaries. K-Theory 4, 55– 88 (1990)
- [Le] Lecomte, F.: Opérations d'Adams en K-Théorie algébrique, Strasbourg, preprint, 1996
- [Lo 1] Loday, J.L.: Homotopie des spaces de concordances (d'aprs F. Waldhausen), Sem. Bourbaki, 1977/78 N. 516, LNM 710, Springer-Verlag, 1979, pp. 187–205
- [Lo 2] Loday, J.L.: Spaces with finitely manynon-trivial homotopy groups, J. of Pure and Applied Algebra 24, 179–202 (1982)
- [Mas] Massey, W.S.: Singular Homology Theory, GTM 70, Springer Verlag, 1980
- [M-S] Milnor, J.W., Stasheff, J.D.: Characteristic Classes, Annals of Mathematics Studies 76, Princeton University Press, 1974
- [N 1] Navarro Aznar, V.: Sur les multiplicités de Schubert locales des faisceaux algébriques cohérents, Comp. Math. 48, 311–326 (1983)
- [N 2] Navarro Aznar, V.: Sur la théorie de Hodge-Deligne, Invent. Math. 90, 11–76 (1987)
- [Ri] Riemenschneider, O.: Characterizing Moishezon spaces by almost positive coherent analytic sheaves, Math. Z. 123, 263–284 (1971)
- [Ro] Rossi, H.: Picard variety of an isolated singular point, Rice Univ. Studies 54, 63–73 (1968)
- [Sch] Schechtman,: On the delooping of Chern character and Adams operations, K-theory, arithmetic and geometry. LNM 1289, Springer-Verlag, 1987, pp. 265–319
- [So] Soulé et als.: Lectures on Arakelov Geometry, Cambridge studies in advanced mathematics 33, Cambridge university Press, 1992
- [Th] Thomason, R.: Algebraic K-theory and étalé cohomology, Ann. Sci. Ec. Norm. Sup. 18, 437–552 (1985)
- [Wal] Waldhausen, F.: Algebraic K-theory of generalized free products, Ann. Math. 108, 135–256 (1978)
- [Wan] Wang X.: Higher-order characteristic classes in arithmetic geometry, Thesis Harvard, 1992